


AD-A236 543




2

HDL-TM-1-1

May 1991

Possible Hosts for Quadruply Ionized  $3d^N$  Transition  
Metal Ions:  $\text{Na}_2\text{TiSiO}_5$ ,  $\text{Y}_2\text{SiBe}_2\text{O}_7$ ,  $\text{Bi}_4\text{X}_3\text{O}_{12}$ , and  $\text{Bi}_{12}\text{XO}_{20}$   
(X = Si, Ge)

by Clyde A. Morrison



U.S. Army Laboratory Command  
Harry Diamond Laboratories  
Adelphi, MD 20783-197

Approved for public release; distribution unlimited.

91-01503



91 6 7 029

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

Citation of manufacturer's or trade names does not constitute an official endorsement or approval of the use thereof.

Destroy this report when it is no longer needed. Do not return it to the originator.

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
<small>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.</small>				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE May 1991		3. REPORT TYPE AND DATES COVERED Interim, 1 Jun 90 to 15 Aug 90
4. TITLE AND SUBTITLE Possible Hosts for Quadruply Ionized $3d^N$ Transition Metal Ions: $\text{Na}_2\text{TiSiO}_5$ , $\text{Y}_2\text{SiBe}_2\text{O}_7$ , $\text{Bi}_4\text{X}_3\text{O}_{12}$ , and $\text{Bi}_{12}\text{XO}_{20}$ ( $X = \text{Si, Ge}$ )			5. FUNDING NUMBERS  DA PR: AH44 PE: 61102H44	
6. AUTHOR(S)  Clyde A. Morrison				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Harry Diamond Laboratories 2800 Powder Mill Road Adelphi, MD 20783-1197			8. PERFORMING ORGANIZATION REPORT NUMBER  HDL-TM-91-1	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Laboratory Command 2800 Powder Mill Road Adelphi, MD 20783-1145			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES AMS code: 611102.H440011 HDL PR: AE1051				
12a. DISTRIBUTION/AVAILABILITY STATEMENT  Approved for public release; distribution unlimited.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words)  This report provides the crystal-field splitting of $\text{Cr}^{4+}$ in the host materials $\text{Na}_2\text{TiSiO}_5$ , $\text{Y}_2\text{SiBe}_2\text{O}_7$ , $\text{Bi}_4\text{X}_3\text{O}_{12}$ , and $\text{Bi}_{12}\text{XO}_{20}$ . The strength of the crystal field is estimated by evaluating the coefficients of a multipole expansion of the crystal field for the Ti, Si, or Ge site and using empirical values of the radial integrals, $\langle r^4 \rangle$ . With the exception of the Ti site in $\text{Na}_2\text{TiSiO}_5$ , the Si or Ge sites ( $D_{2d}$ or $S_4$ symmetry) are, to a certain degree, approximately tetrahedral. In the material $\text{Bi}_{12}\text{XO}_{20}$ the X site (Ge or Si) has the cubic symmetry $T$ . Estimated values of the parameters for the quadruply ionized $3d^N$ electronic configurations are given for $X = \text{Ge}$ . The energy levels are calculated with and without the spin-orbit interaction. The spin-orbit constant is chosen equal to the free-ion value.				
14. SUBJECT TERMS  $\text{Cr}^{4+}$ , $\text{Na}_2\text{TiSiO}_5$ , $\text{Y}_2\text{SiBe}_2\text{O}_7$ , $\text{Bi}_4\text{Ge}_3\text{O}_{12}$ , $\text{Bi}_4\text{Si}_3\text{O}_{12}$ , $\text{Bi}_{12}\text{GeO}_{20}$ , $\text{Bi}_{12}\text{SiO}_{20}$ , $\text{V}^{4+}$ , $\text{Mn}^{4+}$ , $\text{Fe}^{4+}$ , $\text{Co}^{4+}$ , $\text{Ni}^{4+}$ , $\text{Zn}^{4+}$ , $\text{Ga}^{4+}$ , optical spectra			15. NUMBER OF PAGES 56	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	17. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT  UL	

## Contents

	Page
1. Introduction .....	7
2. Theory.....	8
3. Parameters for $\text{Cr}^{4+}$ and Cubic Approximation.....	10
4. Energy Level of $\text{Cr}^{4+}$ in Each Host .....	13
4.1 $\text{Cr}^{4+}:\text{Na}_2\text{TiSiO}_5$ .....	13
4.1.1 Ti Site .....	13
4.1.2 Si Site .....	13
4.2 $\text{Cr}^{4+}:\text{Y}_2\text{SiBe}_2\text{O}_7$ .....	13
4.3 $\text{Cr}^{4+}:\text{Bi}_4\text{X}_3\text{O}_{12}$ ( $\text{X} = \text{Ge}, \text{Si}$ ).....	14
4.3.1 $\text{Bi}_4\text{Si}_3\text{O}_{12}$ .....	14
4.3.2 $\text{Bi}_4\text{Ge}_3\text{O}_{12}$ .....	14
5. Energy Levels of the $\text{X}^{4+}$ Ions in $\text{Bi}_{12}\text{GeO}_{20}$ .....	20
6. Conclusion .....	38
7. References .....	39
Appendix A.—Crystallographic and X-Ray Data .....	41
Distribution .....	51

## Tables

1. Parameters for $\text{Mn}^{4+}$ from Hartree-Fock, free-ion, and $\text{Cs}_2\text{GeF}_6:\text{Mn}^{4+}$ .....	9
2. Free-ion parameters and effective $p_k$ for quadruply ionized ions of the $3d^N$ series derived from $\text{Cs}_2\text{GeF}_6$ experimental.....	11
3. Approximate crystal-field parameters for $\text{Cr}^{4+}$ in the four host materials.....	12
4. Approximate crystal-field parameters for $\text{X}^{4+}$ in the Ge site in $\text{Bi}_{12}\text{GeO}_{20}$ .....	12
5. Energy levels of $\text{Cr}^{4+}$ using the parameters of tables 2 and 3 .....	15
5.1 For Ti site in $\text{Na}_2\text{TiSiO}_5$ , cubic approximation, no spin-orbit .....	15
5.2 For Ti sit in $\text{Na}_2\text{TiSiO}_5$ , $\text{C}_{4v}$ symmetry, no spin-orbit.....	15

## Tables (cont'd)

	Page
5.3 For Si site in $\text{Na}_2\text{TiSiO}_5$ , cubic approximation, no spin-orbit .....	16
5.4 For Si site in $\text{Na}_2\text{TiSiO}_5$ , $D_{2d}$ symmetry, no spin-orbit .....	16
5.5 For Si site in $\text{Y}_2\text{SiBe}_2\text{O}_7$ , cubic approximation, no spin-orbit .....	17
5.6 For Si site in $\text{Y}_2\text{SiBe}_2\text{O}_7$ , $S_4$ symmetry, no spin-orbit .....	17
5.7 For Si site in $\text{Bi}_4\text{Si}_3\text{O}_{12}$ , cubic approximation, no spin-orbit .....	18
5.8 For Si site in $\text{Bi}_4\text{Si}_3\text{O}_{12}$ , $S_4$ symmetry, no spin-orbit .....	18
5.9 For Ge site in $\text{Bi}_4\text{Ge}_3\text{O}_{12}$ , cubic approximation, no spin-orbit .....	19
5.10 For Ge site in $\text{Bi}_4\text{Ge}_3\text{O}_{12}$ , $S_4$ symmetry, no spin-orbit .....	19
 6. Energy levels for $\text{V}^{4+}$ using the parameters of tables 2 and 4, for Ge site in $\text{Bi}_{12}\text{GeO}_{20}$ , $T$ symmetry, no spin orbit .....	 22
7. Energy levels for $\text{V}^{4+}$ using the parameters of tables 2 and 4, for Ge site in $\text{Bi}_{12}\text{GeO}_{20}$ , $T$ symmetry, $\zeta = 250 \text{ cm}^{-1}$ .....	22
8. Energy levels for $\text{Cr}^{4+}$ using the parameters of tables 2 and 4, for Ge site in $\text{Bi}_{12}\text{GeO}_{20}$ , $T$ symmetry, no spin orbit .....	22
9. Energy levels for $\text{Cr}^{4+}$ using the parameters of tables 2 and 4, for Ge site in $\text{Bi}_{12}\text{GeO}_{20}$ , $T$ symmetry, $\zeta = 319 \text{ cm}^{-1}$ .....	23
10. Energy levels for $\text{Mn}^{4+}$ using the parameters of tables 2 and 4, for Ge site in $\text{Bi}_{12}\text{GeO}_{20}$ , $T$ symmetry, no spin orbit .....	24
11. Energy levels for $\text{Mn}^{4+}$ using the parameters of tables 2 and 4, for Ge site in $\text{Bi}_{12}\text{GeO}_{20}$ , $T$ symmetry, $\zeta = 405 \text{ cm}^{-1}$ .....	25
12. Energy levels for $\text{Fe}^{4+}$ using the parameters of tables 2 and 4, for Ge site in $\text{Bi}_{12}\text{GeO}_{12}$ , $T$ symmetry, no spin orbit .....	26
13. Energy levels for $\text{Fe}^{4+}$ using the parameters of tables 2 and 4, for Ge site in $\text{Bi}_{12}\text{GeO}_{20}$ , $T$ symmetry, $\zeta = 513 \text{ cm}^{-1}$ .....	27
14. Energy levels for $\text{Co}^{4+}$ using the parameters of tables 2 and 4, for Ge site in $\text{Bi}_{12}\text{GeO}_{20}$ , $T$ symmetry, no spin orbit .....	29
15. Energy levels for $\text{Co}^{4+}$ using the parameters of tables 2 and 4, for Ge site in $\text{Bi}_{12}\text{GeO}_{20}$ , $T$ symmetry, $\zeta = 654 \text{ cm}^{-1}$ .....	30
16. Energy levels for $\text{Ni}^{4+}$ using the parameters of tables 2 and 4, for Ge site in $\text{Bi}_{12}\text{GeO}_{20}$ , $T$ symmetry, no spin orbit .....	32
17. Energy levels for $\text{Ni}^{4+}$ using the parameters of tables 2 and 4, for Ge site in $\text{Bi}_{12}\text{GeO}_{20}$ , $T$ symmetry, $\zeta = 830 \text{ cm}^{-1}$ .....	33

## Tables (cont'd)

		Page
18.	Energy levels for Cu <sup>4+</sup> using the parameters of tables 2 and 4, for Ge site in Bi <sub>12</sub> GeO <sub>20</sub> , <i>T</i> symmetry, no spin orbit.....	35
19.	Energy levels for Cu <sup>4+</sup> using the parameters of tables 2 and 4, for Ge site in Bi <sub>12</sub> GeO <sub>20</sub> , <i>T</i> symmetry $\zeta = 1008 \text{ cm}^{-1}$ .....	36
20.	Energy levels for Zn <sup>4+</sup> using the parameters of tables 2 and 4, for Ge site in Bi <sub>12</sub> GeO <sub>20</sub> , <i>T</i> symmetry, no spin orbit.....	37
21.	Energy levels for Zn <sup>4+</sup> using the parameters of tables 2 and 4, for Ge site in Bi <sub>12</sub> GeO <sub>20</sub> , <i>T</i> symmetry, $\zeta = 1203 \text{ cm}^{-1}$ .....	37
22.	Energy levels for Ga <sup>4+</sup> using the parameters of tables 2 and 4, for Ge site in Bi <sub>12</sub> GeO <sub>20</sub> , <i>T</i> symmetry, no spin orbit.....	38
23.	Energy levels for Ga <sup>4+</sup> using the parameters of tables 2 and 4, for Ge site in Bi <sub>12</sub> GeO <sub>20</sub> , <i>T</i> symmetry, $\zeta = 1496 \text{ cm}^{-1}$ .....	38



<b>Accession For</b>	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
<b>Availability Codes</b>	
Dist	Avail and/or Special
A-1	

# 1. Introduction

We have selected six compounds as possible host materials for the study of the optical spectra of the quadruply ionized elements with a  $3d^N$  electronic configuration ( $X^{4+}$ ). Previous hosts for which optical spectra of these ions have been investigated have serious drawbacks to the analysis of the experimental data. Frequently the  $X^{4+}$  ion is assumed to enter substitutionally for a triply ionized element of the host, such as  $Mn^{4+}$  in yttrium aluminum garnet (YAG) [1]. In such a case the charge compensation generally perturbs the spectra, giving many more lines than should exist for  $Mn^{4+}$ . Further, the  $Mn^{4+}$  is assumed to go into the  $C_{3i}$  site. If it did, the electric dipole transitions would be forbidden and only magnetic dipole transition would be allowed for the zero phonon lines. Thus, the numerous vibrational assisted electric dipole transitions would be as large or larger than the zero phonon lines. Also, in most of the cases reported, the  $X^{4+}$  ion replaces a quadruply ionized element of the host, but the site symmetry does not allow electric dipole transitions and the spectra must be sorted out from the numerous vibronics. An additional difficulty arises when the symmetry of the site occupied by the transition metal ion is low, such as  $C_2$  symmetry. Then the number of crystal-field parameters is large, thus requiring a large number of levels of the ion to be determined. In our selection of host materials presented here, we have attempted to avoid some of these difficulties.

Each of the host crystals have either  $Si^{4+}$  or  $Ge^{4+}$  as a constituent and we assume that under suitable growth conditions a number of  $X^{4+}$  ions can replace these ions substitutionally. In all six host crystals, the Si or Ge site has approximately tetrahedral symmetry (either  $S_4$  or  $D_{2d}$  symmetry) and in  $Bi_{12}XO_{20}$  ( $X = Ge, Si$ ) the X site has the tetrahedral cubic symmetry,  $T$ . In the crystal  $Na_2TiSiO_5$ , the lighter  $X^{4+}$  ions (V, Cr) might enter the Ti site, which has approximately octahedral symmetry ( $C_{4v}$ ), but the heavier  $X^{4+}$  ions (Ni, Cu) could replace the Si ion, which has approximately tetrahedral symmetry ( $D_{2d}$ ). In all cases the actual symmetry of the site has a maximum of three crystal-field parameters ( $B_{20}$ ,  $B_{40}$ , and  $B_{44}$ ) while in  $Bi_{12}XO_{20}$  the X site has only one crystal-field parameter ( $B_{40}$ ). The symmetry of the Ge and Si sites in all these compounds allows electric dipole transitions so that, in general, the electronic transitions would be stronger than the vibronic lines, and identification of the levels would be easier.

In this report, the detailed x-ray data on the six host materials are given along with references to the growth of these crystals. The x-ray data are used to obtain the crystal-field components,  $A_{kq}$ , and the monopole components of  $A_{kq}$  are used to predict the crystal-field parameters  $B_{kq}$  given by

$$B_{kq} = \rho_k A_{kq} , \quad (1)$$

where the  $\rho_k$  are effective values of  $\langle r^k \rangle$ . The experimental data reported for  $\text{Mn}^{4+}$  in  $\text{Cs}_2\text{GeF}_6$  are used to obtain an approximate set of  $\rho_k$  for the  $X^{4+}$  ions with the  $3d^N$  electronic configuration. Because of the recent interest in the study of  $\text{Cr}^{4+}$  in a crystal-field of tetrahedral symmetry, we have calculated the energy levels of this ion in each of the host materials. In addition, we calculated the energy levels of all the  $X^{4+}$  ions with the  $3d^N$  electronic configuration in the Ge site in  $\text{Bi}_{12}\text{GeO}_{20}$ . This latter calculation is done with and without the spin-orbit interaction.

## 2. Theory

The free ion Hamiltonian for a configuration of  $d^N$  ion is taken as

$$H_{FI} = F^{(2)}g_2 + F^{(4)}g_4 + \zeta \sum_{i=1}^N \vec{l}_i \cdot \vec{s}_i, \quad (2)$$

with

$$g_k = \sum_{i>j} \sum_{q=-k}^k C_{kq}^*(i) C_{kq}(j), \quad (3)$$

and

$$C_{kq} = \sqrt{4\pi/(2k+1)} Y_{kq}. \quad (4)$$

The  $F^{(k)}$  and  $\zeta$  have been calculated with the use of Hartree-Fock wavefunctions [2], and determined by fitting the free-ion spectrum [3] and fitting the spectrum in solids [4]. The matrix elements of  $g_k$ , for all the states of the  $d^N$  configurations, are listed by Nielson and Koster [5], and we use their labels here. Frequently, the Racah parameters  $B$  and  $C$  are used in place of the Slater parameters,  $F^{(k)}$ , and the relations

$$F^{(2)} = 7(7B + C)$$

and

$$F^{(4)} = \frac{63}{5}C \quad (5)$$

can be used to convert from one set of parameters to the other. In general the Slater parameters are considerably reduced when an ion enters a solid. Furthermore, the values for these same parameters obtained by fitting the free ion spectra are less than the corresponding values computed by using Hartree-Fock wavefunctions. As an example of the reduction of the free-ion parameters, we have chosen  $\text{Mn}^{4+}$  in the host  $\text{Cs}_2\text{GeF}_6$  with the parameters for each approximation listed in table 1 (from Morrison [4], p 133).



**Table 1. Parameters for Mn<sup>4+</sup> from Hartree-Fock, free-ion, and Cs<sub>2</sub>GeF<sub>6</sub>:Mn<sup>4+</sup>**

Parameters	Calculated <sup>a</sup> H-F	Experimental <sup>b</sup> free ion	Experimental <sup>c</sup> Cs <sub>2</sub> GeF <sub>6</sub> :Mn <sup>4+</sup>
$F^{(2)} (\text{cm}^{-1})$	101,615	86,939	55,644
$F^{(4)} (\text{cm}^{-1})$	64,078	54,219	49,692
$\zeta (\text{cm}^{-1})$	436	405	380
$\langle r^2 \rangle (\text{\AA}^2)$	0.3568	—	0.8104
$\langle r^4 \rangle (\text{\AA}^4)$	0.4097	—	2.114
$B (\text{cm}^{-1})$	1,347	1,160	572.2
$C (\text{cm}^{-1})$	5,086	4,303	3,944

<sup>a</sup>Morrison [4], p 21; <sup>b</sup>p 14; and <sup>c</sup>p 133.

We see that the reduction of  $B$  in going from the Hartree-Fock value to the value obtained for Mn<sup>4+</sup> in Cs<sub>2</sub>GeF<sub>6</sub> is quite large.

Further, the value  $Dq$  for Mn<sup>4+</sup> in Cs<sub>2</sub>GeF<sub>6</sub> is 2,185 cm<sup>-1</sup> so that  $Dq/B = 3.82$ . Whereas, if the  $Dq$  is computed using  $\langle r^4 \rangle$  from Hartree-Fock values, the crystal-field components,  $A_{40}$ , and equation (1), we have  $Dq = 368.5 \text{ cm}^{-1}$  and  $Dq/B = 0.274$ , which is less than 1/10 the experimental value of  $Dq/B$ . Thus, values of  $Dq/B$  (which is one of the fundamental quantities extracted from comparing experiment when one uses a standard Tanabe-Sugano plot), calculated by Hartree-Fock values greatly underestimate the experimental values. This is not too surprising since the value of  $B$  decreases significantly when the ion enters the solid, whereas the radial wavefunction of the transition metal ion expands, causing an increase of  $\langle r^4 \rangle$ . These two effects cause the ratio  $Dq/B$  to increase dramatically.

From the values given in table 1 we find for Mn<sup>4+</sup> in Cs<sub>2</sub>GeF<sub>6</sub> that  $F^{(2)}/F^{(2)}_{H-F} = 0.5476$ ,  $F^{(4)}/F^{(4)}_{H-F} = 0.7755$ ,  $\langle r^2 \rangle / \langle r^2 \rangle_{H-F} = 2.271$ , and  $\langle r^4 \rangle / \langle r^4 \rangle_{H-F} = 5.159$ . We then assumed for the quadruply ionized 3d<sup>N</sup> ions that  $F^{(2)} = 0.5476 F^{(2)}_{H-F}$ ,  $F^{(4)} = 0.7755 F^{(4)}_{H-F}$ ,  $\rho_2 = 2.271 \langle r^2 \rangle_{H-F}$ , and  $\rho_4 = 5.159 \langle r^4 \rangle_{H-F}$ . These values, along with the free-ion value of  $\zeta$ , are given in table 2.

The crystal-field Hamiltonian for the electronic configuration  $d^N$  appropriate for the hosts here is

$$H_{CEF} = B_{20} \sum_{i=1}^N C_{20}(i) + B_{40} \sum_{i=1}^N C_{40}(i) + B_{44} \sum_{i=1}^N [C_{44}(i) + C_{4-4}(i)] \quad (6)$$

with all  $B_{kq}$  real. In the cubic approximation,  $B_{20} = 0$  and  $B_{44} = (5/14)^{1/2} B_{40}$ . If  $B_{40} < 0$  then we have tetrahedral cubic symmetry and if  $B_{40} > 0$  then the symmetry is octahedral. For both cases,  $B_{40} = 21 Dq$ . A large variety of different parameters are in use for the description of the crystal-field interaction of equation (6); for example, see König and Kremer [6]. The two most frequently quoted are from Ballhausen [7]:

$$\begin{aligned}
B_{20} &= -7Ds , \\
B_{40} &= -21 Dq - 21 Dt , \\
B_{44} &= \frac{3}{2} \sqrt{70} Dq ,
\end{aligned} \tag{7}$$

and from Griffith [8]:

$$\begin{aligned}
B_{20} &= \delta - \mu , \\
B_{40} &= 21 Dq - \left( \delta + \frac{3\mu}{4} \right) \sqrt{\frac{7}{10}} , \\
B_{44} &= 3 \sqrt{\frac{7}{10}} Dq + \left( \delta + \frac{3\mu}{4} \right) \sqrt{\frac{7}{10}} .
\end{aligned} \tag{8}$$

### 3. Parameters for Cr<sup>4+</sup> and Cubic Approximation

The parameters  $\rho_2$  and  $\rho_4$  given in table 2 are used to obtain a set of crystal-field parameters,  $B_{kq}$ , for Cr<sup>4+</sup> in each of the hosts. In obtaining the cubic approximate values, we use the rotational invariant,  $S_k(X)$ , defined as

$$S_k(X) = \left[ \sum_{q=-k}^k X_{kq}^* X_{kq} \right]^{1/2} , \tag{9}$$

where  $X_{kq}$  is a spherical tensor such as the crystal-field components,  $A_{kq}$ , or crystal-field parameters,  $B_{kq}$ . Since for the cubic approximation to the crystal-field given in equation (6) we have  $B_{40}^c = (5/14)^{1/2} B_{44}^c$ , then

$$S_4(B) = \sqrt{\frac{12}{7}} B_{44}^c \tag{10}$$

with

$$S_4(A) = \sqrt{A_{40}^2 + 2A_{44}^2} , \tag{11}$$

and from equation (1) we have

$$B_{40}^c = \sqrt{\frac{7}{12}} \rho_4 S_4(A) , \tag{12}$$

and the sign of  $B_{40}^c$  is taken to be the same as  $A_{40}$ . The result given in equation (12) has been used with the monopole  $A_{kq}$  (see tables A-1.3, -1.4, -2.2, -3.3, and -3.4 in app A) to obtain the cubic approximation to the crystal-field parameters for Cr<sup>4+</sup> in each of the host crystals. The

results are given in table 3 along with the  $Dq/B$  ratio for the cubic approximation ( $B_{40}^C = 21 Dq$ ). In the following text, the results of tables 2 and 3 will be used to calculate energy levels for  $\text{Cr}^{4+}$  in each of the host materials. Both the cubic approximation and the full symmetry of the site are considered. In these calculations we ignore the spin-orbit interaction since it is very small.

Since the Ge site in  $\text{Bi}_{12}\text{GeO}_{20}$  has cubic symmetry, the  $A_{kq}$  from table A-4.4 (see app A) are used to construct the parameters given in table 4.

**Table 2. Free-ion parameters ( $\text{cm}^{-1}$ ) and effective  $\rho_k$  ( $\text{\AA}^k$ ) for quadruply ionized ionic of the  $3d^N$  series derived from  $\text{Cs}_2\text{GeF}_6$  experimental<sup>a</sup>**

Ion	$nd^N$	$F^{(2)}$ $B$	$F^{(4)}$ $C$	$\zeta$	$\rho_2$	$\rho_4$
V	$3d^1$	— —	— —	250	0.9988	3.350
Cr	$3d^2$	52,726 541.6	47,131 3,741	319	0.8421	2.168
Mn	$3d^3$	55,644 572.2	49,692 3,944	405	0.8103	2.114
Fe	$3d^4$	58,465 601.7	52,160 4,140	513	0.6618	1.386
Co	$3d^5$	61,393 632.3	54,736 4,344	654	0.6179	1.172
Ni	$3d^6$	63,611 656.2	56,620 4,494	830	0.5671	0.9673
Cu	$3d^7$	66,073 682.4	58,743 4,662	1,008	0.5185	0.7862
Zn	$3d^8$	68,582 709.1	60,914 4,834	1,203	0.4769	0.6191
Ga	$3d^9$	— —	— —	1,496	0.40878	0.4127

<sup>a</sup> $F^{(2)} = 0.5476 F^{(2)}_{H-F}$ ,  $F^{(4)} = 0.7755 F^{(4)}_{H-F}$ ,  $\zeta = \text{free-ion}$  — except for  $\text{Ga}^{4+}$ ; for  $\text{Ga}^{4+}$ ,  $\zeta = \zeta_{H-F}$  —  $\rho_2 = 2.271 \langle r^2 \rangle_{H-F}$ , and  $\rho_4 = 5.159 \langle r^4 \rangle_{H-F}$ . All the Hartree-Fock data and free-ion data are from Morrison [4], p 21.

**Table 3. Approximate crystal-field parameters (cm<sup>-1</sup>) for Cr<sup>4+</sup> in the four host materials**

No.	Host	Symmetry	$B_{40}^a$	$Dq/B$	$B_{20}$	$B_{40}$	$B_{44}$	Table No.
1	Na <sub>2</sub> TiSiO <sub>5</sub> <sup>b</sup>	$C_{4v}$	39,297	3.46	15,959	39,148	23,610	4.1 and 4.2
2	Na <sub>2</sub> TiSiO <sub>5</sub> <sup>c</sup>	$D_{2d}$	-65,526	5.76	6,135	-68,134	-36,856	4.3 and 4.4
3	Y <sub>2</sub> SiBeO <sub>7</sub>	$S_4$	-59,251	5.21	8,445	-62,421	-32,574	4.5 and 4.6
4	Bi <sub>4</sub> Si <sub>3</sub> O <sub>12</sub>	$S_4$	-68,043	5.98	-3,724	-64,793	-43,236	4.7 and 4.8
5	Bi <sub>4</sub> Ge <sub>3</sub> O <sub>12</sub>	$S_4$	-48,292	4.25	-7,704	-41,352	-33,429	4.9 and 4.10

<sup>a</sup>Cubic approximation  $B_{44} = \sqrt{5/14} B_{40}$ .

<sup>b</sup>Ti site.

<sup>c</sup>Si site.

**Table 4. Approximate crystal-field parameters (cm<sup>-1</sup>) for X<sup>4+</sup> in the Ge site ( $T$  symmetry) in Bi<sub>12</sub>GeO<sub>20</sub>**

Ion	$3d^N$	$B_{40}^a$	$Dq/B$
V	$3d^1$	-84,923	—
Cr	$3d^2$	-54,959	4.83
Mn	$3d^3$	-53,590	4.46
Fe	$3d^4$	-35,135	2.78
Co	$3d^5$	-29,710	2.24
Ni	$3d^6$	-24,521	1.78
Cu	$3d^7$	-19,930	1.39
Zn	$3d^8$	-15,694	1.05
Ga	$3d^9$	-10,462	—

$$^aB_{44} = \sqrt{\frac{5}{14}} B_{40}.$$

## 4. Energy Level of Cr<sup>4+</sup> in Each Host

### 4.1 Cr<sup>4+</sup>:Na<sub>2</sub>TiSiO<sub>5</sub>

#### 4.1.1 Ti Site

The parameters to be used in equations (2) and (6) for the calculation of the energy levels of Cr<sup>4+</sup> in the Ti site from tables 2 and 3 are  $F^{(2)} = 52,726$ ;  $F^{(4)} = 47,131$ ; and  $B_{40}^c = 39,297$  (cm<sup>-1</sup>), with  $B_{44}^c = (5/14)^{1/2} B_{40}^c$ . The results are given in table 5.1. With the same  $F^{(k)}$ , but with  $B_{20} = 15,959$ ;  $B_{40} = 39,148$ ; and  $B_{44} = 23,610$  (cm<sup>-1</sup>) ( $C_{4v}$  symmetry), the results are given in table 5.2. In the cubic symmetry (octahedral) the ground state is  ${}^3T_1(F)$ , which in the full  $C_{4v}$  symmetry splits into a  $\Gamma_5$  ( $E = 0$ ) and  $\Gamma_2$  with a splitting of 6300 cm<sup>-1</sup>. This large splitting is caused by the very large value of  $B_{20}$  since the ratio of  $B_{44}/B_{40}$  is very close to the cubic ratio  $(5/14)^{1/2}$ . The  ${}^1E$  level is split ( $\Gamma_1$  and  $\Gamma_3$  in  $C_{4v}$ ) by approximately 17,000 cm<sup>-1</sup> so that the  $\Gamma_1$  at 6,640 cm<sup>-1</sup> and the  $\Gamma_3$  at 17,899 cm<sup>-1</sup> encompass the  ${}^1T_2$  ( $\Gamma_4$  and  $\Gamma_5$  in  $C_{4v}$ ). However, the largest splitting is the  ${}^3T_1(P)$  at 24,467 cm<sup>-1</sup> which in the  $C_{4v}$  symmetry forms a  $\Gamma_2$  at 18,091 cm<sup>-1</sup> and a  $\Gamma_5$  at 32,363 cm<sup>-1</sup> encompassing the  ${}^3T_2$ , which suffers a comparatively small splitting (~1050 cm<sup>-1</sup>). These results indicate that it would be next to impossible to interpret the experimental data in terms of cubic symmetry, assuming that the approximate values of the parameters are reasonable.

#### 4.1.2 Si Site

The parameter for the cubic approximation is  $B_{40}^c = -65,526$  cm<sup>-1</sup> and for the full  $D_{2d}$  symmetry,  $B_{20} = 6135$ ;  $B_{40} = -68,134$ ; and  $B_{44} = -36,856$  (cm<sup>-1</sup>). The cubic results are given in table 5.3, and the  $D_{2d}$  results are in table 5.4. The  ${}^3A_2$  ground state in cubic symmetry goes over to the  $\Gamma_3$  state. (In the presence of spin-orbit coupling with  $\zeta = 319$  cm<sup>-1</sup>, this state becomes a  $\Gamma_5$  doublet and a  $\Gamma_4$  singlet with the  $\Gamma_5$  lowest and the  $\Gamma_4$  at 0.14 cm<sup>-1</sup>.) The  ${}^1E$  level in cubic symmetry is split into a  $\Gamma_1$  and  $\Gamma_3$  with a splitting of 369 cm<sup>-1</sup>. The  ${}^3T_2$  band is split into a  $\Gamma_5$  and  $\Gamma_4$  band, and the  ${}^3T_1(P)$  is split into a  $\Gamma_2$  and  $\Gamma_5$  band. However, these splittings are not too severe and a first analysis could be done, assuming tetrahedral cubic symmetry.

### 4.2 Cr<sup>4+</sup>:Y<sub>2</sub>SiBe<sub>2</sub>O<sub>7</sub>

The free-ion parameters given in table 2 and the crystal-field parameters given in table 3 are used to calculate the energy levels of Cr<sup>4+</sup> in this host. The calculation for the cubic approximation is given in table 5.5; and for  $S_4$  symmetry, in table 5.6. The  ${}^3A_2$  level in the cubic approximation remains the lowest level in  $S_4$  symmetry despite the

rather large value of  $B_{20}$ . However, the  $^1E$  level is split by  $795\text{ cm}^{-1}$  which might lead to an experimental error in identification using cubic symmetry. Further, the  $^3T_2$  is split by approximately  $3000\text{ cm}^{-1}$  and the  $^3T_1(P)$  is split by approximately  $7500\text{ cm}^{-1}$ , both of which could lead to confusion if the experimental data are interpreted in terms of a cubic approximation. The experimental data might be analyzed in cubic symmetry provided the above deviations are considered.

### 4.3 $\text{Cr}^{4+}:\text{Bi}_4\text{X}_3\text{O}_{12}$ ( $X = \text{Ge}, \text{Si}$ )

#### 4.3.1 $\text{Bi}_4\text{Si}_3\text{O}_{12}$

With the use of the parameters of tables 2 and 3, the energy levels of  $\text{Cr}^{4+}$  in the Si site in this host were calculated, with the results given in tables 5.7 (cubic approximation) and 5.8 ( $S_4$  symmetry). The splitting of the cubic  $^1E$  level by the  $S_4$  symmetry crystal field is only  $6\text{ cm}^{-1}$ , the splitting of the  $^3T_2$  is approximately  $3000\text{ cm}^{-1}$ , and that of the  $^3T_1(P)$  is  $3500\text{ cm}^{-1}$ . The experimental data on  $\text{Cr}^{4+}$  in this host can be crudely analyzed assuming a cubic approximation if one is aware of the possible large splitting of the  $^3T_2$  and  $^3T_1(P)$  states.

#### 4.3.2 $\text{Bi}_4\text{Ge}_3\text{O}_{12}$

The parameters given in tables 2 and 3 are used to calculate the splittings of the levels of  $\text{Cr}^{4+}$  in this host. The results are given in tables 5.9 and 5.10. The splittings in the cubic approximation are smaller than in the previous host. However, the large value of  $B_{20}$  causes a greater splitting of the  $^1E$  ( $66\text{ cm}^{-1}$ ). Also, the large value of  $B_{20}$  causes the  $^3T_1(F)$  to intersperse with the  $^3T_1(P)$ . This mixture of the cubic terms would make the analysis of the experimental data on  $\text{Cr}^{4+}$  in terms of a cubic approximation next to impossible.

**Table 5. Energy levels (cm<sup>-1</sup>) of Cr<sup>4+</sup> using the parameters of tables 2 and 3:  
5.1 For Ti site in Na<sub>2</sub>TiSiO<sub>5</sub>, cubic approximation, no spin-orbit**

Level	I.R. <sup>a</sup>	Energy (cm <sup>-1</sup> )	Free ion state
1	<sup>3</sup> T <sub>1</sub> (F)	0	0.90 <sup>3</sup> F + 0.10 <sup>3</sup> P
2	<sup>1</sup> T <sub>2</sub>	10,979	0.53 <sup>1</sup> D + 0.47 <sup>1</sup> G
3	<sup>1</sup> E	11,075	0.52 <sup>1</sup> G + 0.48 <sup>1</sup> D
4	<sup>3</sup> T <sub>2</sub>	17,527	1.00 <sup>3</sup> F
5	<sup>1</sup> A <sub>1</sub>	23,435	0.71 <sup>1</sup> G + 0.29 <sup>1</sup> S
6	<sup>3</sup> T <sub>1</sub> (P)	24,467	0.90 <sup>3</sup> P + 0.10 <sup>3</sup> F
7	<sup>1</sup> T <sub>2</sub>	29,533	0.53 <sup>1</sup> G + 0.47 <sup>1</sup> D
8	<sup>1</sup> T <sub>1</sub>	31,508	1.00 <sup>1</sup> G
9	<sup>3</sup> A <sub>2</sub>	36,240	1.00 <sup>3</sup> F
10	<sup>1</sup> E	48,150	0.52 <sup>1</sup> D + 0.48 <sup>1</sup> G
11	<sup>1</sup> A <sub>1</sub>	63,701	0.71 <sup>1</sup> S + 0.29 <sup>1</sup> G

<sup>a</sup>Irreducible representation of the O or T<sub>d</sub> group, Mulligan notation.

**5.2 For Ti site in Na<sub>2</sub>TiSiO<sub>5</sub>, C<sub>4v</sub> symmetry, no spin-orbit**

Level	I.R. <sup>a</sup>	Energy (cm <sup>-1</sup> )	Free ion state
1	Γ <sub>5</sub>	0	0.88 <sup>3</sup> F + 0.12 <sup>3</sup> P
2	Γ <sub>2</sub>	6,299	0.97 <sup>3</sup> F + 0.03 <sup>3</sup> P
3	Γ <sub>1</sub>	6,639	0.55 <sup>1</sup> G + 0.41 <sup>1</sup> D + 0.04 <sup>1</sup> S
4	Γ <sub>5</sub>	10,907	0.52 <sup>1</sup> D + 0.48 <sup>1</sup> G
5	Γ <sub>4</sub>	17,751	0.56 <sup>1</sup> D + 0.44 <sup>1</sup> G
6	Γ <sub>3</sub>	17,899	0.51 <sup>1</sup> G + 0.49 <sup>1</sup> D
7	Γ <sub>2</sub>	18,091	0.97 <sup>3</sup> P + 0.03 <sup>3</sup> F
8	Γ <sub>5</sub>	18,601	0.92 <sup>3</sup> F + 0.08 <sup>3</sup> P
9	Γ <sub>4</sub>	19,648	1.00 <sup>3</sup> F
10	Γ <sub>2</sub>	24,595	1.00 <sup>1</sup> G
11	Γ <sub>1</sub>	27,151	0.68 <sup>1</sup> G + 0.22 <sup>1</sup> S + 0.10 <sup>1</sup> D
12	Γ <sub>5</sub>	29,962	0.61 <sup>1</sup> G + 0.39 <sup>1</sup> D
13	Γ <sub>4</sub>	31,719	0.56 <sup>1</sup> G + 0.44 <sup>1</sup> D
14	Γ <sub>5</sub>	32,362	0.80 <sup>3</sup> P + 0.20 <sup>3</sup> F
15	Γ <sub>3</sub>	38,461	1.00 <sup>3</sup> F
16	Γ <sub>5</sub>	40,120	0.91 <sup>1</sup> G + 0.09 <sup>1</sup> D
17	Γ <sub>1</sub>	47,074	0.44 <sup>1</sup> G + 0.39 <sup>1</sup> D + 0.17 <sup>1</sup> S
18	Γ <sub>3</sub>	50,384	0.51 <sup>1</sup> D + 0.49 <sup>1</sup> G
19	Γ <sub>1</sub>	69,763	0.57 <sup>1</sup> S + 0.33 <sup>1</sup> G + 0.10 <sup>1</sup> D

<sup>a</sup>Irreducible representation of the C<sub>4v</sub> group (Bethe notation, G. F. Koster, J. O. Dimmock, R. G. Wheeler, and H. Statz, *Properties of the thirty-two point groups*, MIT Press, Cambridge, MA (1963)).

**C<sub>4v</sub> group selection rules for electric dipole transitions**

**A. Single group**

I.R.	Γ <sub>1</sub>	Γ <sub>2</sub>	Γ <sub>3</sub>	Γ <sub>4</sub>	Γ <sub>5</sub>
Γ <sub>1</sub>	π	0	0	0	σ
Γ <sub>2</sub>	0	π	0	0	σ
Γ <sub>3</sub>	0	0	π	0	σ
Γ <sub>4</sub>	0	0	0	π	σ
Γ <sub>5</sub>	σ	σ	σ	σ	π

**B. Double group**

I.R.	Γ <sub>6</sub>	Γ <sub>7</sub>
Γ <sub>6</sub>	π,σ	σ
Γ <sub>7</sub>	σ	π,σ

Table 5. Energy levels ( $\text{cm}^{-1}$ ) of  $\text{Cr}^{4+}$  using the parameters of tables 2 and 3 (cont'd):  
5.3 For Si site in  $\text{Na}_2\text{TiSiO}_5$ , cubic approximation, no spin-orbit

Level	I.R. <sup>a</sup>	Energy	Free ion state
1	${}^3A_2$	0	$1.00 {}^3F$
2	${}^1E$	11,758	$0.60 {}^1D + 0.40 {}^1G$
3	${}^1A_1$	21,612	$0.76 {}^1G + 0.24 {}^1S$
4	${}^3T_2$	31,202	$1.00 {}^3F$
5	${}^3T_1$	37,307	$0.70 {}^3P + 0.30 {}^3F$
6	${}^1T_2$	42,906	$0.63 {}^1D + 0.37 {}^1G$
7	${}^1T_1$	45,184	$1.00 {}^1G$
8	${}^3T_1$	64,426	$0.70 {}^3F + 0.30 {}^3P$
9	${}^1E$	74,817	$0.60 {}^1G + 0.40 {}^1D$
10	${}^1T_2$	74,872	$0.63 {}^1G + 0.37 {}^1D$
11	${}^1A_1$	92,874	$0.76 {}^1S + 0.24 {}^1G$

<sup>a</sup>Irreducible representation of the  $O$  or  $T_d$  group (Mulligan notation).

5.4 For Si site in  $\text{Na}_2\text{TiSiO}_5$ ,  $D_{2d}$  symmetry, no spin-orbit

Level	I.R. <sup>a</sup>	Energy	Free ion state
1	$\Gamma_3$	0	$1.00 {}^3F$
2	$\Gamma_1$	11,392	$0.58 {}^1D + 0.41 {}^1G + 0.01 {}^1S$
3	$\Gamma_3$	11,760	$0.60 {}^1D + 0.40 {}^1G$
4	$\Gamma_1$	21,961	$0.74 {}^1G + 0.23 {}^1S + 0.02 {}^1D$
5	$\Gamma_4$	29,374	$1.00 {}^3F$
6	$\Gamma_5$	31,960	$0.99 {}^3F + 0.01 {}^3P$
7	$\Gamma_2$	33,585	$0.72 {}^3P + 0.28 {}^3F$
8	$\Gamma_5$	39,196	$0.67 {}^3P + 0.33 {}^3F$
9	$\Gamma_4$	41,092	$0.62 {}^1D + 0.38 {}^1G$
10	$\Gamma_2$	41,385	$1.00 {}^1G$
11	$\Gamma_5$	43,547	$0.60 {}^1D + 0.40 {}^1G$
12	$\Gamma_5$	47,235	$0.96 {}^1G + 0.04 {}^1D$
13	$\Gamma_5$	63,018	$0.68 {}^3F + 0.32 {}^3P$
14	$\Gamma_2$	67,051	$0.72 {}^3F + 0.28 {}^3P$
15	$\Gamma_1$	71,288	$0.59 {}^1G + 0.38 {}^1D + 0.03 {}^1S$
16	$\Gamma_5$	73,418	$0.63 {}^1G + 0.37 {}^1D$
17	$\Gamma_3$	77,517	$0.60 {}^1G + 0.40 {}^1D$
18	$\Gamma_4$	77,560	$0.62 {}^1G + 0.38 {}^1D$
19	$\Gamma_1$	93,496	$0.73 {}^1S + 0.25 {}^1G + 0.01 {}^1D$

<sup>a</sup>Irreducible representations of the  $D_{2d}$  group (G. F. Koster, J. O. Dimmock, R. G. Wheeler, and H. Statz, *Properties of the thirty-two point groups*, MIT Press, Cambridge, MA (1963)).

$D_{2d}$  group selection rules for electric dipole transitions

A. Single group

I.R.	$\Gamma_1$	$\Gamma_2$	$\Gamma_3$	$\Gamma_4$	$\Gamma_5$
$\Gamma_1$	0	0	0	$\pi$	$\sigma$
$\Gamma_2$	0	0	$\pi$	0	$\sigma$
$\Gamma_3$	0	$\pi$	0	0	$\sigma$
$\Gamma_4$	$\pi$	0	0	0	$\sigma$
$\Gamma_5$	$\sigma$	$\sigma$	$\sigma$	$\sigma$	$\pi$

B. Double group

I.R.	$\Gamma_6$	$\Gamma_7$
$\Gamma_6$	$\sigma$	$\pi, \sigma$
$\Gamma_7$	$\pi, \sigma$	$\sigma$



**Table 5. Energy levels (cm<sup>-1</sup>) of Cr<sup>4+</sup> using the parameters of tables 2 and 3 (cont'd):  
5.5 For Si site in Y<sub>2</sub>SiBe<sub>2</sub>O<sub>7</sub>, cubic approximation, no spin orbit**

No.	I.R. <sup>a</sup>	Energy (cm <sup>-1</sup> )	Free ion state
1	<sup>3</sup> A <sub>2</sub>	0	1.00 <sup>3</sup> F
2	<sup>1</sup> E	11,752	0.60 <sup>1</sup> D + 0.40 <sup>1</sup> G
3	<sup>1</sup> A <sub>1</sub>	21,428	0.77 <sup>1</sup> G + 0.23 <sup>1</sup> S
4	<sup>3</sup> T <sub>2</sub>	28,214	1.00 <sup>3</sup> F
5	<sup>3</sup> T <sub>1</sub> (F)	34,270	0.68 <sup>3</sup> P + 0.32 <sup>3</sup> F
6	<sup>1</sup> T <sub>2</sub>	39,907	0.63 <sup>1</sup> D + 0.37 <sup>1</sup> G
7	<sup>1</sup> T <sub>1</sub>	42,195	1.00 <sup>1</sup> G
8	<sup>3</sup> T <sub>1</sub> (F)	58,498	0.68 <sup>3</sup> F + 0.32 <sup>3</sup> P
9	<sup>1</sup> E	68,847	0.60 <sup>1</sup> G + 0.40 <sup>1</sup> D
10	<sup>1</sup> T <sub>2</sub>	68,907	0.63 <sup>1</sup> G + 0.37 <sup>1</sup> D
11	<sup>1</sup> A <sub>1</sub>	87,082	0.77 <sup>1</sup> S + 0.23 <sup>1</sup> G

<sup>a</sup>Irreducible representations of the T<sub>d</sub> or O group (Mulligan notation).

**5.6 For Si site in Y<sub>2</sub>SiBe<sub>2</sub>O<sub>7</sub>, S<sub>4</sub> symmetry, no spin-orbit**

No.	I.R. <sup>a</sup>	Energy (cm <sup>-1</sup> )	Free ion state
1	Γ <sub>2</sub>	0	1.00 <sup>3</sup> F
2	Γ <sub>1</sub>	10,960	0.56 <sup>1</sup> D + 0.42 <sup>1</sup> G + 0.02 <sup>1</sup> S
3	Γ <sub>2</sub>	11,756	0.60 <sup>1</sup> D + 0.40 <sup>1</sup> G
4	Γ <sub>1</sub>	22,179	0.74 <sup>1</sup> G + 0.21 <sup>1</sup> S + 0.05 <sup>1</sup> D
5	Γ <sub>2</sub>	25,955	1.00 <sup>3</sup> F
6	Γ <sub>3,4</sub>	29,044	0.98 <sup>3</sup> F + 0.02 <sup>3</sup> P
7	Γ <sub>1</sub>	29,188	0.71 <sup>3</sup> P + 0.29 <sup>3</sup> F
8	Γ <sub>3,4</sub>	36,886	0.63 <sup>3</sup> P + 0.37 <sup>3</sup> F
9	Γ <sub>1</sub>	36,995	1.00 <sup>1</sup> G
10	Γ <sub>2</sub>	37,668	0.62 <sup>1</sup> D + 0.38 <sup>1</sup> G
11	Γ <sub>3,4</sub>	40,543	0.59 <sup>1</sup> D + 0.41 <sup>1</sup> G
12	Γ <sub>3,4</sub>	45,092	0.94 <sup>1</sup> G + 0.06 <sup>1</sup> D
13	Γ <sub>3,4</sub>	56,644	0.65 <sup>3</sup> F + 0.35 <sup>3</sup> P
14	Γ <sub>1</sub>	61,927	0.71 <sup>3</sup> F + 0.29 <sup>3</sup> P
15	Γ <sub>1</sub>	63,966	0.59 <sup>1</sup> G + 0.37 <sup>1</sup> D + 0.04 <sup>1</sup> S
16	Γ <sub>3,4</sub>	66,965	0.64 <sup>1</sup> G + 0.36 <sup>1</sup> D
17	Γ <sub>2</sub>	72,391	0.60 <sup>1</sup> G + 0.40 <sup>1</sup> D
18	Γ <sub>2</sub>	72,434	0.62 <sup>1</sup> G + 0.38 <sup>1</sup> D
19	Γ <sub>1</sub>	88,092	0.73 <sup>1</sup> S + 0.25 <sup>1</sup> G + 0.02 <sup>1</sup> D

<sup>a</sup>Irreducible representations of the S<sub>4</sub> group, Γ<sub>3,4</sub> = Γ<sub>3</sub> + Γ<sub>4</sub> (Bethe notation, G. F. Koster, J. O. Dimmock, R. G. Wheeler, and H. Statz, Properties of the thirty-two point groups, MIT Press, Cambridge, MA (1963)).

**S<sub>4</sub> group selection rules for electric dipole transitions**

**A. Single group**

I.R.	Γ <sub>1</sub>	Γ <sub>2</sub>	Γ <sub>3</sub>	Γ <sub>4</sub>
Γ <sub>1</sub>	0	π	σ	σ
Γ <sub>2</sub>	π	0	σ	σ
Γ <sub>3</sub>	σ	σ	0	π
Γ <sub>4</sub>	σ	σ	π	0

**B. Double group**

I.R.	Γ <sub>5</sub>	Γ <sub>6</sub>	Γ <sub>7</sub>	Γ <sub>8</sub>
Γ <sub>5</sub>	0	σ	π	σ
Γ <sub>6</sub>	σ	0	σ	π
Γ <sub>7</sub>	π	σ	0	σ
Γ <sub>8</sub>	σ	π	σ	0

**Table 5. Energy levels (cm<sup>-1</sup>) of Cr<sup>4+</sup> using the parameters of tables 2 and 3 (cont'd):  
5.7 For Si site in Bi<sub>4</sub>Si<sub>3</sub>O<sub>12</sub>, cubic approximation, no spin-orbit**

No.	I.R. <sup>a</sup>	Energy (cm <sup>-1</sup> )	Free ion state
1	<sup>3</sup> A <sub>2</sub>	0	1.00 <sup>3</sup> F
2	<sup>1</sup> E	11,760	0.60 <sup>1</sup> D + 0.40 <sup>1</sup> G
3	<sup>1</sup> A <sub>1</sub>	21,678	0.75 <sup>1</sup> G + 0.25 <sup>1</sup> S
4	<sup>3</sup> T <sub>2</sub>	32,401	1.00 <sup>3</sup> F
5	<sup>3</sup> T <sub>1</sub> (P)	38,523	0.70 <sup>3</sup> P + 0.30 <sup>3</sup> F
6	<sup>1</sup> T <sub>2</sub>	44,109	0.63 <sup>1</sup> D + 0.37 <sup>1</sup> G
7	<sup>1</sup> T <sub>1</sub>	46,382	1.00 <sup>1</sup> G
8	<sup>3</sup> T <sub>1</sub> (F)	66,806	0.70 <sup>3</sup> F + 0.30 <sup>3</sup> P
9	<sup>1</sup> E	77,212	0.60 <sup>1</sup> G + 0.40 <sup>1</sup> D
10	<sup>1</sup> T <sub>2</sub>	77,265	0.63 <sup>1</sup> G + 0.37 <sup>1</sup> D
11	<sup>1</sup> A <sub>1</sub>	95,206	0.75 <sup>1</sup> S + 0.25 <sup>1</sup> G

<sup>a</sup>Irreducible representations of the T<sub>d</sub> or O group (Mulligan notation).

**5.8 For Si site in Bi<sub>4</sub>Si<sub>3</sub>O<sub>12</sub>, S<sub>4</sub> symmetry, no spin-orbit**

No.	I.R. <sup>a</sup>	Energy (cm <sup>-1</sup> )	Free ion state
1	Γ <sub>2</sub>	0	1.00 <sup>3</sup> F
2	Γ <sub>1</sub>	11,752	0.60 <sup>1</sup> D + 0.40 <sup>1</sup> G
3	Γ <sub>2</sub>	11,759	0.60 <sup>1</sup> D + 0.40 <sup>1</sup> G
4	Γ <sub>1</sub>	21,680	0.75 <sup>1</sup> G + 0.25 <sup>1</sup> S
5	Γ <sub>3,4</sub>	31,300	1.00 <sup>3</sup> F
6	Γ <sub>2</sub>	34,451	1.00 <sup>3</sup> F
7	Γ <sub>3,4</sub>	37,292	0.71 <sup>3</sup> P + 0.29 <sup>3</sup> F
8	Γ <sub>1</sub>	40,826	0.68 <sup>3</sup> P + 0.32 <sup>3</sup> F
9	Γ <sub>3,4</sub>	43,008	0.62 <sup>1</sup> D + 0.38 <sup>1</sup> G
10	Γ <sub>3,4</sub>	45,129	1.00 <sup>1</sup> G
11	Γ <sub>2</sub>	46,142	0.64 <sup>1</sup> D + 0.36 <sup>1</sup> G
12	Γ <sub>1</sub>	48,761	1.00 <sup>1</sup> G
13	Γ <sub>1</sub>	64,521	0.68 <sup>3</sup> F + 0.32 <sup>3</sup> P
14	Γ <sub>3,4</sub>	67,812	0.71 <sup>3</sup> F + 0.29 <sup>3</sup> P
15	Γ <sub>2</sub>	74,854	0.60 <sup>1</sup> G + 0.40 <sup>1</sup> D
16	Γ <sub>2</sub>	74,921	0.64 <sup>1</sup> G + 0.36 <sup>1</sup> D
17	Γ <sub>3,4</sub>	78,293	0.62 <sup>1</sup> G + 0.38 <sup>1</sup> D
18	Γ <sub>1</sub>	78,772	0.59 <sup>1</sup> G + 0.39 <sup>1</sup> D + 0.03 <sup>1</sup> S
19	Γ <sub>1</sub>	95,721	0.72 <sup>1</sup> S + 0.26 <sup>1</sup> G + 0.01 <sup>1</sup> D

<sup>a</sup>Irreducible representations of the S<sub>4</sub> group, Γ<sub>3,4</sub> = Γ<sub>3</sub> + Γ<sub>4</sub> (Bethe notation, G. F. Koster, J. O. Dimmock, R. G. Wheeler, and H. Statz, *Properties of the thirty-two point groups*, MIT Press, Cambridge, MA (1963)).

Note: See table 5.6 for allowed transitions.

**Table 5. Energy levels (cm<sup>-1</sup>) of Cr<sup>4+</sup> using the parameters of tables 2 and 3 (cont'd):  
5.9 For Ge site in Bi<sub>4</sub>Ge<sub>3</sub>O<sub>12</sub>, cubic approximation, no spin-orbit**

No.	I.R. <sup>a</sup>	Energy (cm <sup>-1</sup> )	Free ion state
1	<sup>3</sup> A <sub>2</sub>	0	1.00 <sup>3</sup> F
2	<sup>1</sup> E	11,738	0.61 <sup>1</sup> D + 0.39 <sup>1</sup> G
3	<sup>1</sup> A <sub>1</sub>	21,015	0.80 <sup>1</sup> G + 0.20 <sup>1</sup> S
4	<sup>3</sup> T <sub>2</sub>	22,996	1.00 <sup>3</sup> F
5	<sup>3</sup> T <sub>1</sub> (P)	28,931	0.65 <sup>3</sup> P + 0.35 <sup>3</sup> F
6	<sup>1</sup> T <sub>2</sub>	34,662	0.65 <sup>1</sup> D + 0.35 <sup>1</sup> G
7	<sup>1</sup> T <sub>1</sub>	36,977	1.00 <sup>1</sup> G
8	<sup>3</sup> T <sub>1</sub> (F)	48,182	0.65 <sup>3</sup> F + 0.35 <sup>3</sup> P
9	<sup>1</sup> E	58,424	0.61 <sup>1</sup> G + 0.39 <sup>1</sup> D
10	<sup>1</sup> T <sub>2</sub>	58,497	0.65 <sup>1</sup> G + 0.35 <sup>1</sup> D
11	<sup>1</sup> A <sub>1</sub>	77,058	0.80 <sup>1</sup> S + 0.20 <sup>1</sup> G

<sup>a</sup>Irreducible representations of the T<sub>d</sub> or O group (Mulligan notation).

**5.10 For Ge site in Bi<sub>4</sub>Ge<sub>3</sub>O<sub>12</sub>, S<sub>4</sub> symmetry, no spin-orbit**

No.	I.R. <sup>a</sup>	Energy (cm <sup>-1</sup> )	Free ion state
1	Γ <sub>2</sub>	0	1.00 <sup>3</sup> F
2	Γ <sub>1</sub>	11,663	0.60 <sup>1</sup> D + 0.40 <sup>1</sup> G
3	Γ <sub>3,4</sub>	11,729	0.62 <sup>1</sup> D + 0.38 <sup>1</sup> G
4	Γ <sub>3,4</sub>	20,531	1.00 <sup>3</sup> F
5	Γ <sub>1</sub>	21,004	0.80 <sup>1</sup> G + 0.19 <sup>1</sup> S + 0.01 <sup>1</sup> D
6	Γ <sub>3,4</sub>	26,162	0.67 <sup>3</sup> P + 0.33 <sup>3</sup> F
7	Γ <sub>2</sub>	26,636	1.00 <sup>3</sup> F
8	Γ <sub>3,4</sub>	32,162	0.61 <sup>1</sup> D + 0.39 <sup>1</sup> G
9	Γ <sub>1</sub>	32,936	0.51 <sup>3</sup> F + 0.49 <sup>3</sup> P
10	Γ <sub>3,4</sub>	34,160	0.97 <sup>1</sup> G + 0.03 <sup>1</sup> D
11	Γ <sub>2</sub>	38,213	0.69 <sup>1</sup> D + 0.31 <sup>1</sup> G
12	Γ <sub>1</sub>	41,547	1.00 <sup>1</sup> G
13	Γ <sub>1</sub>	43,409	0.51 <sup>3</sup> P + 0.49 <sup>3</sup> F
14	Γ <sub>3,4</sub>	49,514	0.68 <sup>3</sup> F + 0.32 <sup>3</sup> P
15	Γ <sub>2</sub>	53,095	0.62 <sup>1</sup> G + 0.38 <sup>1</sup> D
16	Γ <sub>2</sub>	53,248	0.69 <sup>1</sup> G + 0.31 <sup>1</sup> D
17	Γ <sub>1</sub>	59,816	0.55 <sup>1</sup> G + 0.34 <sup>1</sup> D + 0.11 <sup>1</sup> S
18	Γ <sub>3,4</sub>	59,912	0.64 <sup>1</sup> G + 0.36 <sup>1</sup> D
19	Γ <sub>1</sub>	78,626	0.69 <sup>1</sup> S + 0.26 <sup>1</sup> G + 0.05 <sup>1</sup> D

<sup>a</sup>Irreducible representations of the S<sub>4</sub> group, Γ<sub>3,4</sub> = Γ<sub>3</sub> + Γ<sub>4</sub> (Bethe notation, G. F. Koster, J. O. Dimmock, R. G. Wheeler, and H. Statz, *Properties of the thirty-two point groups*, MIT Press, Cambridge, MA (1963)).

Note: See table 5.6 for allowed transitions.

## 5. Energy Levels of the $X^{4+}$ Ions in $\text{Bi}_{12}\text{GeO}_{20}$

Because of the technological importance of the compounds  $\text{Bi}_{12}\text{XO}_{20}$  ( $X = \text{Ge}, \text{Si}$ ) [9,10], we have decided to calculate the energy levels of the entire  $3d^N$  series of  $X^{4+}$  ions assuming these ions replace the Ge site substitutionally. Also, the Ge (Si) site in these compounds has the unique distinction of being of the cubic group  $T$ , and the optical spectra of these ions should be quite interesting. The parameters used in the calculations are given in tables 2 and 4, and the results are given in tables 6 to 23. For each ion, the energy levels are calculated with the spin-orbit interaction "turned off," followed by a similar calculation with the spin-orbit parameter values given in table 2. The optical spectra of  $\text{Cr}^{4+}$  in  $\text{Bi}_{12}\text{GeO}_{20}$  have been reported by Wardzynski et al [11], who give  $B_{40} = -17,220 \text{ cm}^{-1}$  and  $B = 428.8 \text{ cm}^{-1}$ , giving the parameter  $Dq/B = 1.91$ , whereas our estimated value,  $Dq/B = 4.83$ , is much larger. This would indicate that our estimate of  $Dq$  is too large or  $B$  is too small. However, their reported energy levels correspond more closely to the levels of  $\text{Fe}^{4+}$  (tables 12 and 13) calculated here.

The labeling of the levels is the irreducible representation of the  $T$  groups in the Bethe notation [12]. The correspondence with the  $T_d$  group in Mulligan notation is as follows:  $A_1$  and  $A_2$  correspond to  $\Gamma_1$ ,  $E$  becomes  $\Gamma_2 + \Gamma_3$  (still a doublet), and  $T_1$  and  $T_2$  correspond to the triplet  $\Gamma_4$ . The double group  $\Gamma_6$  and  $\Gamma_7$  ( $T_d$ ) corresponds to  $\Gamma_5$  ( $T$ ), and the quartet  $\Gamma_8$  ( $T_d$ ) becomes  $\Gamma_6 + \Gamma_7$  ( $T$ ), which is two degenerate doublets. For  $d$  electrons the difference between the  $T$  and  $T_d$  groups is insignificant, but for  $f$  electrons there is a distinct difference. In the absence of the spin-orbit interaction, the energy levels can be obtained from a Tanabe-Sugano plot for ions with the  $d^N$  configurations provided the complementary rules are observed for tetrahedral symmetry. That is, for the configuration  $d^N$  in tetrahedral symmetry, we use the  $d^{10-N}$  Tanabe-Sugano plot. (These standard plots are for octahedral symmetry.) The inclusion of the spin-orbit interaction has very little effect on the  $\text{V}^{4+}$ ,  $\text{Cr}^{4+}$ , and  $\text{Mn}^{4+}$  ions. However, in the  $\text{Fe}^{4+}$  ion (table 13) the  $\Gamma_1$  ground state in the absence of spin-orbit (40 percent  $^1I$ ) remains  $\Gamma_1$  but is 67 percent  $^5D$  with 11 percent  $^1I$ . This mixture could be changed considerably by different values of  $Dq/B$ . The spin-orbit interaction causes the  $^6S$  ground state of  $\text{Co}^{4+}$  with  $\zeta = 0$  to split into a  $\Gamma_5$  level at  $E = 0$  and a four-fold degenerate level  $\Gamma_{6,7}$  to split at  $0.04 \text{ cm}^{-1}$  (table 15).

For the ion  $\text{Ni}^{4+}$  the  $\Gamma_{2,3}$  level of the ground state with  $\zeta = 0$  is split into 5 levels ( $0 \leq E \leq 176 \text{ cm}^{-1}$ ) by the spin-orbit interaction, as shown in tables 16 and 17. Also the spin-orbit mixes different spin states beginning with the  $\Gamma_4$  levels at  $11,142 \text{ cm}^{-1}$ . The  $\text{Cu}^{4+}$  ion energy levels given in table 19 show that the first excited state is mixed by the spin-orbit interaction (98 percent  $^4F + 1$  percent  $^2G$ ) with the higher energy

levels becoming strongly mixed. Similarly, for the energy levels of  $\text{Zn}^{4+}$ , table 21, and  $\text{Ga}^{4+}$ , table 23, the spin-orbit interaction has a reasonably strong effect on the composition of the wave functions for the higher energy levels. This mixture of different total spin raises the spin forbidden transition rules usually employed in the interpretation of  $3d^N$  electronic configuration optical absorption data.

The only ion with experimental data other than  $\text{Cr}^{4+}$  that we were able to find was  $\text{Fe}^{4+}$  [13], and we have not investigated this ionization state here.

The interstitial site mentioned [14] for  $\text{Bi}_{12}\text{GeO}_{20}$  was investigated. The statement concerning the size of this site (6b in the crystallographic data) is correct. In the host material  $\text{Bi}_{12}\text{GeO}_{20}$ , the nearest neighbors at this site have four Bi ions at 2.86 Å while the Bi site has eight oxygen ions ranging from 2.08 to 3.47 Å and the Ge ion has four oxygen ions at 1.72 Å. However, because the nearest neighbors are  $\text{Bi}^{3+}$  it seems that it would be difficult to trap a positive ion at this position. On the other hand, this site might act as an electron trapping center in this crystal (also the silicon counterpart). Nevertheless, the monopole crystal-field components,  $A_{kq}$  ( $\text{cm}^{-1}/\text{\AA}^k$ ), were calculated, with  $A_{20} = -31,713$ ;  $A_{22} = 19,514$ ;  $A_{40} = 4,521$ ;  $A_{42} = 7,866$ ; and  $A_{44} = -613$ , and the site has  $D_2$  symmetry. If this site had  $D_{2d}$  symmetry then  $A_{22}$  and  $A_{42}$  would vanish, which appears to be rather a gross assumption considering the relative magnitude of these components.

A second possible interstitial site not mentioned by Wardzynski and Szymczak [14] is the site (12d) which is located at (0, 0, 1/4) and has  $C_2$  symmetry. This site has for nearest neighbors two Bi ions at 1.92 Å, two O ions at 2.09 Å, and one Ge ion at 2.54 Å. The monopole crystal-field components  $A_{kq}$  ( $\text{cm}^{-1}/\text{\AA}^k$ ) are as follows:  $A_{20} = 12,856$ ;  $A_{22} = 47,509 + i15,341$ ;  $A_{40} = -7,415$ ;  $A_{42} = -111 + i6,203$ ; and  $A_{44} = -11,806 - i3,334$ . The two-fold fields are slightly larger (absolute magnitude) than the previous interstitial site but this site has, in addition, a large  $A_{10}$  component ( $48,637 \text{ cm}^{-1}/\text{\AA}$ ) which is equivalent to a constant electric field in the  $z$  direction. In view of these factors, this site does not appear to be a possible electron trapping site for electrons, but further work would be necessary to determine if a doubly ionized transition-metal ion might occupy it.

**Table 6. Energy levels (cm<sup>-1</sup>) for V<sup>4+</sup> using the parameters of tables 2 and 4, for Ge site in Bi<sub>12</sub>GeO<sub>20</sub>, *T* symmetry, no spin orbit**

No.	I.R. <sup>a</sup>	Energy (cm <sup>-1</sup> )	Free ion state
1	$\Gamma_{2,3}$	0	1.00 <sup>2</sup> D
2	$\Gamma_4$	40,439	1.00 <sup>2</sup> D

<sup>a</sup>Irreducible representations of the *T* group,  $\Gamma_{2,3} = \Gamma_2 + \Gamma_3$  (Bethe notation, G. F. Koster, J. O. Dimmock, R. G. Wheeler, and H. Statz, *Properties of the thirty-two point groups*, MIT Press, Cambridge, MA (1963)).

**Table 7. Energy levels (cm<sup>-1</sup>) for V<sup>4+</sup> using the parameters of tables 2 and 4, for Ge site in Bi<sub>12</sub>GeO<sub>20</sub>, *T* symmetry,  $\zeta = 250$  cm<sup>-1</sup>**

No.	I.R. <sup>a</sup>	Energy (cm <sup>-1</sup> )	Free ion state
1	$\Gamma_{6,7}$	0	1.00 <sup>2</sup> D
2	$\Gamma_{6,7}$	40,319	1.00 <sup>2</sup> D
3	$\Gamma_5$	40,691	1.00 <sup>2</sup> D

<sup>a</sup>Irreducible representations of the double *T* (cubic) group (G. F. Koster, J. O. Dimmock, R. G. Wheeler, and H. Statz, *Properties of the thirty-two point groups*, MIT Press, Cambridge, MA (1963)).

**Table 8. Energy levels (cm<sup>-1</sup>) for Cr<sup>4+</sup> using the parameters of tables 2 and 4, for Ge site in Bi<sub>12</sub>GeO<sub>20</sub>, *T* symmetry, no spin orbit**

No.	I.R. <sup>a</sup>	Energy (cm <sup>-1</sup> )	Free ion state
1	$\Gamma_1$	0	1.00 <sup>3</sup> F
2	$\Gamma_{2,3}$	11,748	0.61 <sup>1</sup> D + 0.39 <sup>1</sup> G
3	$\Gamma_1$	21,282	0.78 <sup>1</sup> G + 0.22 <sup>1</sup> S
4	$\Gamma_4$	26,170	1.00 <sup>3</sup> F
5	$\Gamma_4$	32,186	0.67 <sup>3</sup> P + 0.33 <sup>3</sup> F
6	$\Gamma_4$	37,854	0.64 <sup>1</sup> D + 0.36 <sup>1</sup> G
7	$\Gamma_4$	40,152	1.00 <sup>1</sup> G
8	$\Gamma_4$	54,451	0.67 <sup>3</sup> F + 0.33 <sup>3</sup> P
9	$\Gamma_{2,3}$	64,764	0.61 <sup>1</sup> G + 0.39 <sup>1</sup> D
10	$\Gamma_4$	64,829	0.64 <sup>1</sup> G + 0.36 <sup>1</sup> D
11	$\Gamma_1$	83,140	0.78 <sup>1</sup> S + 0.22 <sup>1</sup> G

<sup>a</sup>Irreducible representations of the *T* (cubic) group,  $\Gamma_{2,3} = \Gamma_2 + \Gamma_3$  (Bethe notation, G. F. Koster, J. O. Dimmock, R. G. Wheeler, and H. Statz, *Properties of the thirty-two point groups*, MIT Press, Cambridge, MA (1963)).

**Table 9.** Energy levels ( $\text{cm}^{-1}$ ) for  $\text{Cr}^{4+}$  using the parameters of tables 2 and 4, for Ge site in  $\text{Bi}_{12}\text{GeO}_{20}$ ,  $T$  symmetry,  $\zeta = 319 \text{ cm}^{-1}$

No.	I.R. <sup>a</sup>	Energy ( $\text{cm}^{-1}$ )	Free ion state		
1	$\Gamma_4$	0	1.00 $^3F$		
2	$\Gamma_{2,3}$	11,739	0.61 $^1D$	+	0.39 $^1S$
3	$\Gamma_1$	21,273	0.78 $^1G$	+	0.22 $^1S$
4	$\Gamma_1$	26,021	1.00 $^3F$		
5	$\Gamma_4$	26,096	1.00 $^3F$		
6	$\Gamma_{2,3}$	26,249	1.00 $^3F$		
7	$\Gamma_4$	26,251	1.00 $^3F$		
8	$\Gamma_1$	32,149	0.68 $^3P$	+	0.31 $^3F$
9	$\Gamma_4$	32,169	0.68 $^3P$	+	0.32 $^3F$
10	$\Gamma_4$	32,223	0.66 $^3P$	+	0.34 $^3F$
11	$\Gamma_{2,3}$	32,239	0.66 $^3P$	+	0.34 $^3F$
12	$\Gamma_4$	37,861	0.64 $^1D$	+	0.36 $^1S$
13	$\Gamma_4$	40,164	1.00 $^1G$		
14	$\Gamma_{2,3}$	54,360	0.66 $^3F$	+	0.34 $^3P$
15	$\Gamma_4$	54,365	0.66 $^3F$	+	0.34 $^3P$
16	$\Gamma_4$	54,577	0.68 $^3F$	+	0.32 $^3P$
17	$\Gamma_1$	54,674	0.69 $^3F$	+	0.31 $^3P$
18	$\Gamma_{2,3}$	64,784	0.61 $^1G$	+	0.39 $^1D$
19	$\Gamma_4$	64,849	0.64 $^1G$	+	0.36 $^1D$
20	$\Gamma_1$	83,165	0.78 $^1S$	+	0.22 $^1S$

<sup>a</sup>Irreducible representations of the  $T$  group,  $\Gamma_{2,3} = \Gamma_2 + \Gamma_3$  (degenerate doublets) (G. F. Koster, J. O. Dimmock, R. G. Wheeler, and H. Statz, *Properties of the thirty-two point groups*, MIT Press, Cambridge, MA (1963)).

**Table 10.** Energy levels ( $\text{cm}^{-1}$ ) for  $\text{Mn}^{4+}$  using the parameters of tables 2 and 4, for Ge site in  $\text{Bi}_{12}\text{GeO}_{20}$ ,  $T$  symmetry, no spin orbit

No.	I.R. <sup>a</sup>	Energy ( $\text{cm}^{-1}$ )	Free ion state		
1	$\Gamma_{2,3}$	0	0.40 $^2G$	+	0.32 $^2H$ + 0.15 $^2D_1$
2	$\Gamma_4$	8,233	0.88 $^4F$	+	0.12 $^4P$
3	$\Gamma_4$	22,675	0.56 $^2H$	+	0.44 $^2G$
4	$\Gamma_4$	23,652	0.51 $^2G$	+	0.24 $^2D_2$ + 0.16 $^2H$
5	$\Gamma_4$	26,721	0.54 $^2P$	+	0.27 $^2F$ + 0.19 $^2H$
6	$\Gamma_4$	32,415	1.00 $^4F$		
7	$\Gamma_4$	37,865	0.37 $^2H$	+	0.29 $^2F$ + 0.24 $^2D_1$
8	$\Gamma_4$	39,661	0.88 $^4P$	+	0.12 $^4F$
9	$\Gamma_1$	46,535	1.00 $^2G$		
10	$\Gamma_4$	48,689	0.60 $^2H$	+	0.25 $^2G$ + 0.14 $^2P$
11	$\Gamma_4$	49,882	0.43 $^2G$	+	0.30 $^2D_2$ + 0.15 $^2H$
12	$\Gamma_{2,3}$	50,480	0.50 $^2H$	+	0.43 $^2D_2$ + 0.06 $^2D_1$
13	$\Gamma_4$	53,810	0.55 $^2F$	+	0.35 $^2H$ + 0.06 $^2G$
14	$\Gamma_4$	54,639	0.40 $^2D_2$	+	0.40 $^2F$ + 0.18 $^2H$
15	$\Gamma_1$	57,979	1.00 $^2F$		
16	$\Gamma_1$	57,980	1.00 $^2F$		
17	$\Gamma_{2,3}$	68,057	0.51 $^2D_1$	+	0.44 $^2G$ + 0.05 $^2H$
18	$\Gamma_4$	75,290	0.30 $^2H$	+	0.28 $^2P$ + 0.24 $^2G$
19	$\Gamma_{2,3}$	79,031	0.43 $^2D_2$	+	0.28 $^2D_1$ + 0.15 $^2G$
20	$\Gamma_4$	89,511	0.68 $^2D_1$	+	0.16 $^2F$ + 0.14 $^2H$

<sup>a</sup>Irreducible representations of the  $T$  group,  $\Gamma_{2,3} = \Gamma_2 + \Gamma_3$  (Bethe notation, G. F. Koster, J. O. Dimmock, R. G. Wheeler, and H. Statz, *Properties of the thirty-two point groups*, MIT Press, Cambridge, MA (1963)).



**Table 11. Energy levels (cm<sup>-1</sup>) for Mn<sup>4+</sup> using the parameters of tables 2 and 4, for Ge site in Bi<sub>12</sub>GeO<sub>20</sub>, *T* symmetry,  $\zeta = 405$  cm<sup>-1</sup>**

No.	I.R. <sup>a</sup>	Energy (cm <sup>-1</sup> )	Free ion state				
1	$\Gamma_{6,7}$	0	0.40 <sup>2</sup> G	+	0.32 <sup>2</sup> H	+	0.15 <sup>2</sup> D1
2	$\Gamma_5$	8,034	0.88 <sup>4</sup> F	+	0.12 <sup>4</sup> P		
3	$\Gamma_{6,7}$	8,055	0.88 <sup>4</sup> F	+	0.12 <sup>4</sup> P		
4	$\Gamma_{6,7}$	8,431	0.88 <sup>4</sup> F	+	0.12 <sup>4</sup> P		
5	$\Gamma_5$	8,674	0.87 <sup>4</sup> F	+	0.13 <sup>4</sup> P		
6	$\Gamma_5$	22,548	0.55 <sup>2</sup> H	+	0.44 <sup>2</sup> G		
7	$\Gamma_{6,7}$	22,771	0.55 <sup>2</sup> H	+	0.44 <sup>2</sup> G	+	0.01 <sup>2</sup> D2
8	$\Gamma_5$	23,534	0.52 <sup>2</sup> G	+	0.24 <sup>2</sup> D2	+	0.15 <sup>2</sup> H
9	$\Gamma_{6,7}$	23,739	0.50 <sup>2</sup> G	+	0.23 <sup>2</sup> D2	+	0.18 <sup>2</sup> H
10	$\Gamma_5$	26,549	0.54 <sup>2</sup> P	+	0.27 <sup>2</sup> F	+	0.19 <sup>2</sup> H
11	$\Gamma_{6,7}$	26,864	0.54 <sup>2</sup> P	+	0.27 <sup>2</sup> F	+	0.19 <sup>2</sup> H
12	$\Gamma_5$	32,310	1.00 <sup>4</sup> F				
13	$\Gamma_{6,7}$	32,426	0.99 <sup>4</sup> F				
14	$\Gamma_{6,7}$	32,566	1.00 <sup>4</sup> F				
15	$\Gamma_5$	32,574	1.00 <sup>4</sup> F				
16	$\Gamma_{6,7}$	37,747	0.35 <sup>2</sup> H	+	0.29 <sup>2</sup> F	+	0.24 <sup>2</sup> D1
17	$\Gamma_5$	38,097	0.37 <sup>2</sup> H	+	0.26 <sup>2</sup> F	+	0.24 <sup>2</sup> D1
18	$\Gamma_5$	39,481	0.87 <sup>4</sup> P	+	0.13 <sup>4</sup> F		
19	$\Gamma_{6,7}$	39,642	0.87 <sup>4</sup> P	+	0.12 <sup>4</sup> F		
20	$\Gamma_{6,7}$	39,879	0.87 <sup>4</sup> P	+	0.12 <sup>4</sup> F	+	0.01 <sup>2</sup> H
21	$\Gamma_5$	39,925	0.85 <sup>4</sup> P	+	0.11 <sup>4</sup> F	+	0.02 <sup>2</sup> H
22	$\Gamma_5$	46,589	1.00 <sup>2</sup> G				
23	$\Gamma_5$	48,613	0.63 <sup>2</sup> H	+	0.23 <sup>2</sup> G	+	0.14 <sup>2</sup> P
24	$\Gamma_{6,7}$	48,803	0.58 <sup>2</sup> H	+	0.26 <sup>2</sup> G	+	0.15 <sup>2</sup> P
25	$\Gamma_{6,7}$	49,850	0.41 <sup>2</sup> G	+	0.29 <sup>2</sup> D2	+	0.17 <sup>2</sup> H
26	$\Gamma_5$	49,965	0.42 <sup>2</sup> G	+	0.33 <sup>2</sup> D2	+	0.13 <sup>2</sup> H
27	$\Gamma_{6,7}$	50,568	0.49 <sup>2</sup> H	+	0.42 <sup>2</sup> D2	+	0.06 <sup>2</sup> D1
28	$\Gamma_{6,7}$	53,819	0.54 <sup>2</sup> F	+	0.36 <sup>2</sup> H	+	0.06 <sup>2</sup> G
29	$\Gamma_5$	53,962	0.56 <sup>2</sup> F	+	0.32 <sup>2</sup> H	+	0.08 <sup>2</sup> G
30	$\Gamma_{6,7}$	54,674	0.41 <sup>2</sup> D2	+	0.39 <sup>2</sup> F	+	0.18 <sup>2</sup> H
31	$\Gamma_5$	54,746	0.43 <sup>2</sup> F	+	0.37 <sup>2</sup> D2	+	0.18 <sup>2</sup> H
32	$\Gamma_{6,7}$	58,029	0.99 <sup>4</sup> F				
33	$\Gamma_5$	58,038	1.00 <sup>2</sup> F				
34	$\Gamma_{6,7}$	68,125	0.51 <sup>2</sup> D1	+	0.44 <sup>2</sup> G	+	0.05 <sup>2</sup> H
35	$\Gamma_5$	75,294	0.31 <sup>2</sup> H	+	0.27 <sup>2</sup> P	+	0.25 <sup>2</sup> G
36	$\Gamma_{6,7}$	75,380	0.30 <sup>2</sup> H	+	0.28 <sup>2</sup> P	+	0.24 <sup>2</sup> G
37	$\Gamma_{6,7}$	79,098	0.43 <sup>2</sup> D2	+	0.28 <sup>2</sup> D1	+	0.15 <sup>2</sup> G
38	$\Gamma_5$	89,441	0.68 <sup>2</sup> D1	+	0.15 <sup>2</sup> F	+	0.15 <sup>2</sup> H
39	$\Gamma_{6,7}$	89,657	0.68 <sup>2</sup> D1	+	0.16 <sup>2</sup> F	+	0.14 <sup>2</sup> H

<sup>a</sup>Irreducible representations of the double T (cubic) group (G. F. Koster, J. O. Dimmock, R. G. Wheeler, and H. Statz, *Properties of the thirty-two point groups*, MIT Press, Cambridge, MA (1963)).

Table 12. Energy levels ( $\text{cm}^{-1}$ ) for  $\text{Fe}^{4+}$  using the parameters of tables 2 and 4, for Ge site in  $\text{Bi}_{12}\text{GeO}_{20}$ ,  $T$  symmetry, no spin orbit

No.	I.R. <sup>a</sup>	Energy ( $\text{cm}^{-1}$ )	Free ion state					
1	$\Gamma_1$	0	0.40 $^1I$	+	0.30 $^1G_1$	+	0.22 $^1G_2$	
2	$\Gamma_4$	45	1.00 $^5D$					
3	$\Gamma_4$	5,601	0.47 $^3H$	+	0.25 $^3F_2$	+	0.15 $^3P_1$	
4	$\Gamma_4$	9,139	0.40 $^3G$	+	0.27 $^3H$	+	0.17 $^3F_1$	
5	$\Gamma_4$	14,290	0.62 $^1I$	+	0.21 $^1G_1$	+	0.17 $^1G_2$	
6	$\Gamma_{2,3}$	16,776	1.00 $^5D$					
7	$\Gamma_4$	19,938	0.67 $^3H$	+	0.28 $^3F_2$	+	0.04 $^3F_1$	
8	$\Gamma_4$	20,554	0.52 $^3H$	+	0.43 $^3G$	+	0.04 $^3F_2$	
9	$\Gamma_4$	21,631	0.36 $^1I$	+	0.20 $^1F$	+	0.19 $^1D_2$	
10	$\Gamma_{2,3}$	22,245	0.52 $^3G$	+	0.30 $^3H$	+	0.18 $^3D$	
11	$\Gamma_4$	22,983	0.50 $^3G$	+	0.37 $^3P_2$	+	0.05 $^3P_1$	
12	$\Gamma_4$	24,961	0.53 $^3D$	+	0.43 $^3F_2$	+	0.03 $^3G$	
13	$\Gamma_4$	29,791	0.62 $^1I$	+	0.17 $^1G_2$	+	0.11 $^1D_2$	
14	$\Gamma_{2,3}$	30,722	0.51 $^1G_2$	+	0.25 $^1I$	+	0.17 $^1G_1$	
15	$\Gamma_1$	31,331	0.48 $^1I$	+	0.28 $^1S_2$	+	0.16 $^1G_2$	
16	$\Gamma_4$	33,577	0.52 $^3F_1$	+	0.48 $^3F_2$			
17	$\Gamma_4$	34,226	0.43 $^1G_2$	+	0.32 $^1F$	+	0.21 $^1I$	
18	$\Gamma_1$	34,661	0.54 $^1F$	+	0.46 $^1I$			
19	$\Gamma_4$	35,518	0.52 $^3H$	+	0.28 $^3F_2$	+	0.10 $^3P_1$	
20	$\Gamma_4$	35,647	0.30 $^1D_2$	+	0.27 $^1F$	+	0.25 $^1I$	
21	$\Gamma_{2,3}$	36,622	0.68 $^3H$	+	0.30 $^3G$	+	0.02 $^3D$	
22	$\Gamma_4$	38,743	0.38 $^3F_2$	+	0.24 $^3D$	+	0.20 $^3H$	
23	$\Gamma_{2,3}$	38,750	1.00 $^3G$					
24	$\Gamma_4$	38,832	0.40 $^3P_2$	+	0.35 $^3G$	+	0.14 $^3F_1$	
25	$\Gamma_{2,3}$	41,856	0.80 $^3D$	+	0.18 $^3G$	+	0.02 $^3H$	
26	$\Gamma_{2,3}$	43,396	0.45 $^1D_2$	+	0.40 $^1G_1$	+	0.12 $^1D_1$	
27	$\Gamma_4$	44,946	0.49 $^1I$	+	0.23 $^1G_1$	+	0.21 $^1D_2$	
28	$\Gamma_{2,3}$	46,774	0.55 $^1I$	+	0.35 $^1G_2$	+	0.03 $^1D_1$	
29	$\Gamma_4$	48,037	0.52 $^3F_2$	+	0.48 $^3F_1$			
30	$\Gamma_1$	48,084	0.54 $^1I$	+	0.46 $^1F$			
31	$\Gamma_4$	50,506	0.47 $^1G_2$	+	0.27 $^1F$	+	0.09 $^1D_1$	
32	$\Gamma_1$	50,732	0.45 $^1G_2$	+	0.42 $^1S_2$	+	0.09 $^1S_1$	
33	$\Gamma_4$	50,904	0.65 $^1F$	+	0.13 $^1G_2$	+	0.13 $^1I$	
34	$\Gamma_4$	55,156	0.70 $^3P_1$	+	0.13 $^3F_2$	+	0.08 $^3P_2$	
35	$\Gamma_4$	55,479	0.74 $^3F_1$	+	0.11 $^3F_2$	+	0.10 $^3D$	
36	$\Gamma_4$	61,956	0.61 $^1G_1$	+	0.17 $^1I$	+	0.09 $^1D_2$	
37	$\Gamma_4$	61,971	0.65 $^3F_1$	+	0.14 $^3H$	+	0.12 $^3G$	
38	$\Gamma_{2,3}$	64,355	0.50 $^1D_1$	+	0.36 $^1D_2$	+	0.07 $^1G_2$	
39	$\Gamma_1$	64,403	0.50 $^1G_1$	+	0.18 $^1S_2$	+	0.14 $^1G_2$	
40	$\Gamma_4$	64,766	0.67 $^1G_1$	+	0.26 $^1G_2$	+	0.05 $^1I$	
41	$\Gamma_{2,3}$	73,171	0.34 $^1D_1$	+	0.33 $^1G_1$	+	0.16 $^1I$	
42	$\Gamma_4$	79,955	0.69 $^1D_1$	+	0.14 $^1F$	+	0.09 $^1D_2$	
43	$\Gamma_1$	100,394	0.72 $^1S_1$	+	0.18 $^1G_1$	+	0.06 $^1S_2$	

<sup>a</sup>Irreducible representations of the  $T$  (cubic) group,  $\Gamma_{2,3} = \Gamma_2 + \Gamma_3$  (Bethe notation, G. F. Koster, J. O. Dimmock, R. G. Wheeler, and H. Statz, *Properties of the thirty-two point groups*, MIT Press, Cambridge, MA (1963)).

Table 13. Energy levels (cm<sup>-1</sup>) for Fe<sup>4+</sup> using the parameters of tables 2 and 4, for Ge site in Bi<sub>12</sub>GeO<sub>20</sub>, *T* symmetry,  $\zeta = 513$  cm<sup>-1</sup>

No.	I.R. <sup>a</sup>	Energy (cm <sup>-1</sup> )	Free ion state				
1	Γ <sub>1</sub>	0	0.67 <sup>5</sup> D	+	0.11 <sup>1</sup> I	+	0.08 <sup>1</sup> G1
2	Γ <sub>4</sub>	159	0.99 <sup>5</sup> D				
3	Γ <sub>4</sub>	208	0.99 <sup>5</sup> D				
4	Γ <sub>1</sub>	354	0.31 <sup>5</sup> D	+	0.27 <sup>1</sup> I	+	0.20 <sup>1</sup> G1
5	Γ <sub>2,3</sub>	579	0.99 <sup>5</sup> D				
6	Γ <sub>4</sub>	606	1.00 <sup>5</sup> D				
7	Γ <sub>4</sub>	868	1.00 <sup>5</sup> D				
8	Γ <sub>4</sub>	5,962	0.47 <sup>3</sup> H	+	0.24 <sup>3</sup> F2	+	0.15 <sup>3</sup> P1
9	Γ <sub>1</sub>	6,007	0.44 <sup>3</sup> H	+	0.22 <sup>3</sup> F2	+	0.15 <sup>3</sup> P1
10	Γ <sub>2,3</sub>	6,196	0.47 <sup>3</sup> H	+	0.26 <sup>3</sup> F2	+	0.14 <sup>3</sup> P1
11	Γ <sub>4</sub>	6,253	0.47 <sup>3</sup> H	+	0.26 <sup>3</sup> F2	+	0.14 <sup>3</sup> P1
12	Γ <sub>1</sub>	9,214	0.44 <sup>3</sup> G	+	0.24 <sup>3</sup> H	+	0.17 <sup>3</sup> F1
13	Γ <sub>4</sub>	9,430	0.42 <sup>3</sup> G	+	0.26 <sup>3</sup> H	+	0.17 <sup>3</sup> F1
14	Γ <sub>4</sub>	9,760	0.38 <sup>3</sup> G	+	0.29 <sup>3</sup> H	+	0.16 <sup>3</sup> F1
15	Γ <sub>2,3</sub>	9,838	0.38 <sup>3</sup> G	+	0.29 <sup>3</sup> H	+	0.16 <sup>3</sup> F1
16	Γ <sub>4</sub>	14,690	0.60 <sup>1</sup> I	+	0.21 <sup>1</sup> G1	+	0.17 <sup>1</sup> G2
17	Γ <sub>1</sub>	17,147	0.97 <sup>5</sup> D	+	0.01 <sup>3</sup> H	+	0.01 <sup>3</sup> F2
18	Γ <sub>4</sub>	17,164	0.98 <sup>5</sup> D	+	0.01 <sup>3</sup> H	+	0.01 <sup>3</sup> F2
19	Γ <sub>2,3</sub>	17,174	0.98 <sup>5</sup> D	+	0.01 <sup>3</sup> H	+	0.01 <sup>3</sup> F2
20	Γ <sub>4</sub>	17,183	0.99 <sup>5</sup> D				
21	Γ <sub>1</sub>	17,193	0.99 <sup>5</sup> D				
22	Γ <sub>2,3</sub>	20,302	0.65 <sup>3</sup> H	+	0.22 <sup>3</sup> F2	+	0.09 <sup>3</sup> G
23	Γ <sub>4</sub>	20,387	0.66 <sup>3</sup> H	+	0.25 <sup>3</sup> F2	+	0.04 <sup>3</sup> G
24	Γ <sub>1</sub>	20,410	0.65 <sup>3</sup> H	+	0.29 <sup>3</sup> F2	+	0.04 <sup>3</sup> F1
25	Γ <sub>4</sub>	20,433	0.64 <sup>3</sup> H	+	0.26 <sup>3</sup> F2	+	0.04 <sup>3</sup> G
26	Γ <sub>4</sub>	21,055	0.51 <sup>3</sup> H	+	0.41 <sup>3</sup> G	+	0.07 <sup>3</sup> F2
27	Γ <sub>1</sub>	21,061	0.55 <sup>3</sup> H	+	0.41 <sup>3</sup> G	+	0.03 <sup>3</sup> F2
28	Γ <sub>4</sub>	21,064	0.53 <sup>3</sup> H	+	0.37 <sup>3</sup> G	+	0.07 <sup>3</sup> F2
29	Γ <sub>2,3</sub>	21,174	0.53 <sup>3</sup> H	+	0.36 <sup>3</sup> G	+	0.10 <sup>3</sup> F2
30	Γ <sub>4</sub>	21,813	0.27 <sup>1</sup> I	+	0.15 <sup>1</sup> F	+	0.15 <sup>1</sup> D2
31	Γ <sub>4</sub>	22,725	0.51 <sup>3</sup> G	+	0.31 <sup>3</sup> H	+	0.18 <sup>3</sup> D
32	Γ <sub>4</sub>	22,884	0.43 <sup>3</sup> G	+	0.25 <sup>3</sup> H	+	0.15 <sup>3</sup> D
33	Γ <sub>1</sub>	23,358	0.46 <sup>3</sup> G	+	0.37 <sup>3</sup> P2	+	0.06 <sup>3</sup> H
34	Γ <sub>4</sub>	23,413	0.48 <sup>3</sup> G	+	0.37 <sup>3</sup> P2	+	0.05 <sup>3</sup> H
35	Γ <sub>2,3</sub>	23,506	0.51 <sup>3</sup> G	+	0.36 <sup>3</sup> P2	+	0.05 <sup>3</sup> P1
36	Γ <sub>4</sub>	23,545	0.50 <sup>3</sup> G	+	0.36 <sup>3</sup> P2	+	0.05 <sup>3</sup> P1
37	Γ <sub>1</sub>	25,473	0.54 <sup>3</sup> D	+	0.43 <sup>3</sup> F2	+	0.02 <sup>3</sup> G
38	Γ <sub>2,3</sub>	25,479	0.51 <sup>3</sup> D	+	0.43 <sup>3</sup> F2	+	0.05 <sup>3</sup> G
39	Γ <sub>4</sub>	25,484	0.52 <sup>3</sup> D	+	0.43 <sup>3</sup> F2	+	0.05 <sup>3</sup> G
40	Γ <sub>4</sub>	25,535	0.51 <sup>3</sup> D	+	0.42 <sup>3</sup> F2	+	0.03 <sup>3</sup> G
41	Γ <sub>4</sub>	30,220	0.61 <sup>1</sup> I	+	0.17 <sup>1</sup> G2	+	0.11 <sup>1</sup> D2
42	Γ <sub>2,3</sub>	31,154	0.50 <sup>1</sup> G2	+	0.25 <sup>1</sup> I	+	0.17 <sup>1</sup> G1
43	Γ <sub>1</sub>	31,733	0.47 <sup>1</sup> I	+	0.27 <sup>1</sup> S2	+	0.16 <sup>1</sup> G2
44	Γ <sub>4</sub>	34,103	0.52 <sup>3</sup> F1	+	0.47 <sup>3</sup> F2		
45	Γ <sub>4</sub>	34,531	0.39 <sup>1</sup> G2	+	0.30 <sup>1</sup> F	+	0.19 <sup>1</sup> I
46	Γ <sub>1</sub>	35,089	0.53 <sup>1</sup> F	+	0.45 <sup>1</sup> I	+	0.01 <sup>3</sup> F2
47	Γ <sub>4</sub>	35,924	0.34 <sup>3</sup> H	+	0.17 <sup>3</sup> F2	+	0.11 <sup>1</sup> F
48	Γ <sub>2,3</sub>	35,984	0.51 <sup>3</sup> H	+	0.29 <sup>3</sup> F2	+	0.10 <sup>3</sup> P1
49	Γ <sub>4</sub>	36,060	0.19 <sup>1</sup> D2	+	0.19 <sup>3</sup> H	+	0.15 <sup>1</sup> F

Table 13. Energy levels ( $\text{cm}^{-1}$ ) for  $\text{Fe}^{4+}$  using the parameters of tables 2 and 4, for Ge site in  $\text{Bi}_{12}\text{GeO}_{20}$ ,  $T$  symmetry,  $\zeta = 513 \text{ cm}^{-1}$  (cont'd)

No.	I.R. <sup>a</sup>	Energy (cm <sup>-1</sup> )	Free ion state				
50	Γ <sub>1</sub>	36,127	0.52 <sup>3</sup> H	+	0.27 <sup>3</sup> F <sub>2</sub>	+	0.10 <sup>3</sup> P <sub>1</sub>
51	Γ <sub>4</sub>	36,138	0.49 <sup>3</sup> H	+	0.26 <sup>3</sup> F <sub>2</sub>	+	0.09 <sup>3</sup> P <sub>1</sub>
52	Γ <sub>4</sub>	37,135	0.67 <sup>3</sup> H	+	0.29 <sup>3</sup> G	+	0.02 <sup>3</sup> D
53	Γ <sub>4</sub>	37,170	0.66 <sup>3</sup> H	+	0.28 <sup>3</sup> G	+	0.02 <sup>3</sup> D
54	Γ <sub>4</sub>	39,070	0.31 <sup>3</sup> F <sub>2</sub>	+	0.18 <sup>3</sup> D	+	0.17 <sup>3</sup> H
55	Γ <sub>2,3</sub>	39,088	0.35 <sup>3</sup> F <sub>2</sub>	+	0.21 <sup>3</sup> D	+	0.20 <sup>3</sup> H
56	Γ <sub>4</sub>	39,172	0.64 <sup>3</sup> G	+	0.12 <sup>3</sup> P <sub>2</sub>	+	0.07 <sup>3</sup> F <sub>2</sub>
57	Γ <sub>1</sub>	39,179	0.38 <sup>3</sup> G	+	0.37 <sup>3</sup> P <sub>2</sub>	+	0.15 <sup>3</sup> F <sub>1</sub>
58	Γ <sub>4</sub>	39,358	0.25 <sup>3</sup> P <sub>2</sub>	+	0.24 <sup>3</sup> G	+	0.16 <sup>3</sup> F <sub>2</sub>
59	Γ <sub>4</sub>	39,377	0.63 <sup>3</sup> G	+	0.20 <sup>3</sup> P <sub>2</sub>	+	0.07 <sup>3</sup> F <sub>1</sub>
60	Γ <sub>2,3</sub>	39,418	0.38 <sup>3</sup> P <sub>2</sub>	+	0.30 <sup>3</sup> G	+	0.14 <sup>3</sup> F <sub>1</sub>
61	Γ <sub>4</sub>	39,483	0.24 <sup>3</sup> F <sub>2</sub>	+	0.17 <sup>3</sup> G	+	0.17 <sup>3</sup> P <sub>2</sub>
62	Γ <sub>1</sub>	39,590	0.38 <sup>3</sup> F <sub>2</sub>	+	0.25 <sup>3</sup> D	+	0.19 <sup>3</sup> H
63	Γ <sub>4</sub>	42,346	0.79 <sup>3</sup> D	+	0.18 <sup>3</sup> G	+	0.02 <sup>3</sup> H
64	Γ <sub>4</sub>	42,348	0.80 <sup>3</sup> D	+	0.18 <sup>3</sup> G	+	0.02 <sup>3</sup> H
65	Γ <sub>2,3</sub>	43,871	0.44 <sup>1</sup> D <sub>2</sub>	+	0.39 <sup>1</sup> G <sub>1</sub>	+	0.12 <sup>1</sup> D <sub>1</sub>
66	Γ <sub>4</sub>	45,448	0.49 <sup>1</sup> I	+	0.23 <sup>1</sup> G <sub>1</sub>	+	0.21 <sup>1</sup> D <sub>2</sub>
67	Γ <sub>2,3</sub>	47,259	0.55 <sup>1</sup> I	+	0.35 <sup>1</sup> G <sub>2</sub>	+	0.04 <sup>1</sup> D <sub>1</sub>
68	Γ <sub>4</sub>	48,532	0.52 <sup>3</sup> F <sub>2</sub>	+	0.48 <sup>3</sup> F <sub>1</sub>		
69	Γ <sub>1</sub>	48,545	0.54 <sup>1</sup> I	+	0.46 <sup>1</sup> F		
70	Γ <sub>4</sub>	50,955	0.47 <sup>1</sup> G <sub>2</sub>	+	0.27 <sup>1</sup> F	+	0.09 <sup>1</sup> D <sub>1</sub>
71	Γ <sub>1</sub>	51,112	0.44 <sup>1</sup> G <sub>2</sub>	+	0.41 <sup>1</sup> S <sub>2</sub>	+	0.08 <sup>1</sup> S <sub>1</sub>
72	Γ <sub>4</sub>	51,359	0.64 <sup>1</sup> F	+	0.13 <sup>1</sup> G <sub>2</sub>	+	0.13 <sup>1</sup> I
73	Γ <sub>2,3</sub>	55,463	0.67 <sup>3</sup> P <sub>1</sub>	+	0.14 <sup>3</sup> F <sub>2</sub>	+	0.07 <sup>3</sup> H
74	Γ <sub>4</sub>	55,471	0.68 <sup>3</sup> P <sub>1</sub>	+	0.14 <sup>3</sup> F <sub>2</sub>	+	0.07 <sup>3</sup> H
75	Γ <sub>1</sub>	55,822	0.74 <sup>3</sup> F <sub>1</sub>	+	0.10 <sup>3</sup> F <sub>2</sub>	+	0.08 <sup>3</sup> D
76	Γ <sub>4</sub>	55,839	0.66 <sup>3</sup> P <sub>1</sub>	+	0.13 <sup>3</sup> F <sub>2</sub>	+	0.09 <sup>3</sup> P <sub>2</sub>
77	Γ <sub>4</sub>	55,926	0.72 <sup>3</sup> F <sub>1</sub>	+	0.11 <sup>3</sup> F <sub>2</sub>	+	0.09 <sup>3</sup> D
78	Γ <sub>2,3</sub>	56,104	0.70 <sup>3</sup> F <sub>1</sub>	+	0.11 <sup>3</sup> F <sub>2</sub>	+	0.10 <sup>3</sup> D
79	Γ <sub>4</sub>	56,129	0.69 <sup>3</sup> F <sub>1</sub>	+	0.11 <sup>3</sup> F <sub>2</sub>	+	0.10 <sup>3</sup> D
80	Γ <sub>1</sub>	56,168	0.67 <sup>3</sup> P <sub>1</sub>	+	0.12 <sup>3</sup> F <sub>2</sub>	+	0.09 <sup>3</sup> P <sub>2</sub>
81	Γ <sub>4</sub>	62,162	0.37 <sup>1</sup> G <sub>1</sub>	+	0.26 <sup>3</sup> F <sub>1</sub>	+	0.11 <sup>1</sup> I
82	Γ <sub>1</sub>	62,219	0.66 <sup>3</sup> F <sub>1</sub>	+	0.14 <sup>3</sup> H	+	0.11 <sup>3</sup> G
83	Γ <sub>4</sub>	62,378	0.66 <sup>3</sup> F <sub>1</sub>	+	0.14 <sup>3</sup> H	+	0.12 <sup>3</sup> G
84	Γ <sub>2,3</sub>	62,590	0.64 <sup>3</sup> F <sub>1</sub>	+	0.14 <sup>3</sup> H	+	0.13 <sup>3</sup> G
85	Γ <sub>4</sub>	62,963	0.39 <sup>3</sup> F <sub>1</sub>	+	0.24 <sup>1</sup> G <sub>1</sub>	+	0.09 <sup>3</sup> H
86	Γ <sub>2,3</sub>	64,886	0.49 <sup>1</sup> D <sub>1</sub>	+	0.35 <sup>1</sup> D <sub>2</sub>	+	0.07 <sup>1</sup> G <sub>2</sub>
87	Γ <sub>1</sub>	64,921	0.50 <sup>1</sup> G <sub>1</sub>	+	0.18 <sup>1</sup> S <sub>2</sub>	+	0.14 <sup>1</sup> G <sub>2</sub>
88	Γ <sub>4</sub>	65,279	0.67 <sup>1</sup> G <sub>1</sub>	+	0.26 <sup>1</sup> G <sub>2</sub>	+	0.04 <sup>1</sup> I
89	Γ <sub>2,3</sub>	73,700	0.33 <sup>1</sup> D <sub>1</sub>	+	0.33 <sup>1</sup> G <sub>1</sub>	+	0.16 <sup>1</sup> I
90	Γ <sub>4</sub>	80,458	0.69 <sup>1</sup> D <sub>1</sub>	+	0.14 <sup>1</sup> F	+	0.09 <sup>1</sup> D <sub>2</sub>
91	Γ <sub>1</sub>	100,908	0.72 <sup>1</sup> S <sub>1</sub>	+	0.18 <sup>1</sup> G <sub>1</sub>	+	0.06 <sup>1</sup> S <sub>2</sub>

<sup>a</sup>Irreducible representations of the  $T$  (cubic) group,  $\Gamma_{2,3} = \Gamma_2 + \Gamma_3$  (G. F. Koster, J. O. Dimmock, R. G. Wheeler, and H. Statz, *Properties of the thirty-two point groups*, MIT Press, Cambridge, MA (1963)).

**Table 14. Energy levels (cm<sup>-1</sup>) for Co<sup>4+</sup> using the parameters of tables 2 and 4, for Ge site in Bi<sub>12</sub>GeO<sub>20</sub>, *T* symmetry, no spin orbit**

No.	I.R. <sup>a</sup>	Energy (cm <sup>-1</sup> )	Free ion state				
1	Γ <sub>1</sub>	0	1.00 <sup>6</sup> S				
2	Γ <sub>4</sub>	17,382	0.63 <sup>4</sup> G	+	0.33 <sup>4</sup> P	+	0.03 <sup>4</sup> F
3	Γ <sub>4</sub>	21,309	0.35 <sup>2</sup> I	+	0.26 <sup>2</sup> H	+	0.17 <sup>2</sup> F1
4	Γ <sub>4</sub>	21,316	0.50 <sup>4</sup> G	+	0.26 <sup>4</sup> F	+	0.24 <sup>4</sup> D
5	Γ <sub>4</sub>	28,043	1.00 <sup>4</sup> G				
6	Γ <sub>4</sub>	30,284	0.57 <sup>4</sup> D	+	0.41 <sup>4</sup> G	+	0.02 <sup>4</sup> F
7	Γ <sub>1</sub>	31,705	0.59 <sup>2</sup> I	+	0.37 <sup>2</sup> F1	+	0.04 <sup>2</sup> F2
8	Γ <sub>4</sub>	31,834	0.63 <sup>2</sup> I	+	0.23 <sup>2</sup> H	+	0.12 <sup>2</sup> F1
9	Γ <sub>2,3</sub>	32,470	1.00 <sup>4</sup> D				
10	Γ <sub>4</sub>	33,261	0.52 <sup>2</sup> I	+	0.22 <sup>2</sup> F1	+	0.10 <sup>2</sup> H
11	Γ <sub>2,3</sub>	35,005	0.43 <sup>2</sup> I	+	0.25 <sup>2</sup> H	+	0.12 <sup>2</sup> D3
12	Γ <sub>4</sub>	38,873	0.37 <sup>2</sup> G2	+	0.20 <sup>2</sup> F2	+	0.18 <sup>2</sup> F1
13	Γ <sub>1</sub>	40,501	0.71 <sup>2</sup> I	+	0.15 <sup>2</sup> G2	+	0.10 <sup>2</sup> G1
14	Γ <sub>4</sub>	40,547	0.32 <sup>2</sup> G2	+	0.24 <sup>2</sup> F2	+	0.18 <sup>2</sup> I
15	Γ <sub>4</sub>	41,108	0.45 <sup>4</sup> P	+	0.43 <sup>4</sup> F	+	0.11 <sup>4</sup> G
16	Γ <sub>2,3</sub>	43,278	0.45 <sup>2</sup> I	+	0.29 <sup>2</sup> D3	+	0.11 <sup>2</sup> H
17	Γ <sub>1</sub>	43,831	0.51 <sup>2</sup> G2	+	0.28 <sup>2</sup> I	+	0.12 <sup>2</sup> G1
18	Γ <sub>4</sub>	44,300	0.45 <sup>2</sup> I	+	0.39 <sup>2</sup> D3	+	0.06 <sup>2</sup> D1
19	Γ <sub>1</sub>	44,320	1.00 <sup>4</sup> F				
20	Γ <sub>2,3</sub>	47,294	0.73 <sup>2</sup> G2	+	0.12 <sup>2</sup> D3	+	0.05 <sup>2</sup> D1
21	Γ <sub>4</sub>	48,545	0.46 <sup>2</sup> F2	+	0.27 <sup>2</sup> G2	+	0.12 <sup>2</sup> I
22	Γ <sub>4</sub>	48,708	0.53 <sup>4</sup> F	+	0.26 <sup>4</sup> G	+	0.21 <sup>4</sup> P
23	Γ <sub>1</sub>	49,327	0.75 <sup>2</sup> F2	+	0.18 <sup>2</sup> I	+	0.07 <sup>2</sup> F1
24	Γ <sub>4</sub>	49,402	0.52 <sup>2</sup> F2	+	0.37 <sup>2</sup> G2	+	0.07 <sup>2</sup> I
25	Γ <sub>4</sub>	51,107	0.68 <sup>2</sup> H	+	0.31 <sup>2</sup> F1		
26	Γ <sub>1</sub>	52,702	0.71 <sup>2</sup> S	+	0.24 <sup>2</sup> G2	+	0.04 <sup>2</sup> G1
27	Γ <sub>4</sub>	53,233	0.71 <sup>4</sup> F	+	0.19 <sup>4</sup> D	+	0.10 <sup>4</sup> G
28	Γ <sub>2,3</sub>	55,188	0.44 <sup>2</sup> H	+	0.30 <sup>2</sup> D3	+	0.24 <sup>2</sup> D2
29	Γ <sub>4</sub>	56,796	0.41 <sup>2</sup> H	+	0.20 <sup>2</sup> F1	+	0.16 <sup>2</sup> F2
30	Γ <sub>4</sub>	58,217	0.34 <sup>2</sup> F1	+	0.22 <sup>2</sup> H	+	0.13 <sup>2</sup> G2
31	Γ <sub>4</sub>	61,526	0.45 <sup>2</sup> H	+	0.27 <sup>2</sup> I	+	0.18 <sup>2</sup> F1
32	Γ <sub>1</sub>	61,634	0.56 <sup>2</sup> F1	+	0.23 <sup>2</sup> I	+	0.21 <sup>2</sup> F2
33	Γ <sub>2,3</sub>	64,912	0.78 <sup>2</sup> G1	+	0.08 <sup>2</sup> H	+	0.07 <sup>2</sup> D1
34	Γ <sub>4</sub>	67,311	0.49 <sup>2</sup> D2	+	0.17 <sup>2</sup> G1	+	0.13 <sup>2</sup> I
35	Γ <sub>2,3</sub>	68,375	0.60 <sup>2</sup> D2	+	0.10 <sup>2</sup> H	+	0.10 <sup>2</sup> G2
36	Γ <sub>4</sub>	69,487	0.26 <sup>2</sup> D3	+	0.22 <sup>2</sup> H	+	0.18 <sup>2</sup> F1
37	Γ <sub>4</sub>	70,166	0.80 <sup>2</sup> G1	+	0.07 <sup>2</sup> P	+	0.06 <sup>2</sup> F2
38	Γ <sub>4</sub>	73,792	0.52 <sup>2</sup> G1	+	0.13 <sup>2</sup> F2	+	0.12 <sup>2</sup> G2
39	Γ <sub>1</sub>	75,794	0.74 <sup>2</sup> G1	+	0.16 <sup>2</sup> S	+	0.10 <sup>2</sup> G2
40	Γ <sub>4</sub>	80,636	0.87 <sup>2</sup> P	+	0.04 <sup>2</sup> F2	+	0.04 <sup>2</sup> G2
41	Γ <sub>4</sub>	94,481	0.73 <sup>2</sup> D1	+	0.07 <sup>2</sup> H	+	0.05 <sup>2</sup> D3
42	Γ <sub>2,3</sub>	94,533	0.73 <sup>2</sup> D1	+	0.10 <sup>2</sup> G1	+	0.10 <sup>2</sup> D3

<sup>a</sup>Irreducible representations of the *T* (cubic) group,  $\Gamma_{2,3} = \Gamma_2 + \Gamma_3$  (Bethe notation, G. F. Koster, J. O. Dimmock, R. G. Wheeler, and H. Statz, *Properties of the thirty-two point groups*, MIT Press, Cambridge, MA (1963)).

Table 15. Energy levels (cm<sup>-1</sup>) for Co<sup>4+</sup> using the parameters of tables 2 and 4, for Ge site in Bi<sub>12</sub>GeO<sub>20</sub>, *T* symmetry,  $\zeta = 654$  cm<sup>-1</sup>

No.	I.R. <sup>a</sup>	Energy (cm <sup>-1</sup> )	Free ion state			
1	$\Gamma_{6,7}$	0	1.00 <sup>6</sup> S			
2	$\Gamma_{6,7}$	0.04	1.00 <sup>6</sup> S			
3	$\Gamma_5$	17,231	0.63 <sup>4</sup> G	+	0.33 <sup>4</sup> P	+ 0.04 <sup>4</sup> F
4	$\Gamma_{6,7}$	17,247	0.61 <sup>4</sup> G	+	0.32 <sup>4</sup> P	+ 0.04 <sup>4</sup> F
5	$\Gamma_5$	17,313	0.60 <sup>4</sup> G	+	0.32 <sup>4</sup> P	+ 0.03 <sup>4</sup> F
6	$\Gamma_{6,7}$	17,410	0.61 <sup>4</sup> G	+	0.33 <sup>4</sup> P	+ 0.03 <sup>4</sup> F
7	$\Gamma_{6,7}$	20,762	0.22 <sup>4</sup> G	+	0.19 <sup>2</sup> I	+ 0.14 <sup>2</sup> H
8	$\Gamma_5$	21,241	0.43 <sup>4</sup> G	+	0.26 <sup>4</sup> D	+ 0.25 <sup>4</sup> F
9	$\Gamma_5$	21,350	0.53 <sup>4</sup> G	+	0.26 <sup>4</sup> F	+ 0.21 <sup>4</sup> D
10	$\Gamma_{6,7}$	21,374	0.53 <sup>4</sup> G	+	0.26 <sup>4</sup> F	+ 0.21 <sup>4</sup> D
11	$\Gamma_{6,7}$	21,689	0.29 <sup>4</sup> G	+	0.16 <sup>4</sup> F	+ 0.16 <sup>4</sup> D
12	$\Gamma_5$	22,297	0.34 <sup>2</sup> I	+	0.25 <sup>2</sup> H	+ 0.13 <sup>2</sup> F1
13	$\Gamma_{6,7}$	27,957	0.97 <sup>4</sup> G	+	0.01 <sup>2</sup> I	+ 0.01 <sup>2</sup> F1
14	$\Gamma_5$	27,995	0.98 <sup>4</sup> G	+	0.01 <sup>2</sup> F1	
15	$\Gamma_5$	28,019	0.98 <sup>4</sup> G	+	0.01 <sup>2</sup> I	+ 0.01 <sup>2</sup> H
16	$\Gamma_{6,7}$	28,052	0.99 <sup>4</sup> G			
17	$\Gamma_5$	29,754	0.46 <sup>4</sup> D	+	0.40 <sup>4</sup> G	+ 0.06 <sup>2</sup> I
18	$\Gamma_{6,7}$	30,194	0.54 <sup>4</sup> D	+	0.43 <sup>4</sup> G	+ 0.02 <sup>4</sup> F
19	$\Gamma_{6,7}$	30,416	0.59 <sup>4</sup> D	+	0.37 <sup>4</sup> G	+ 0.02 <sup>4</sup> F
20	$\Gamma_5$	30,468	0.60 <sup>4</sup> D	+	0.37 <sup>4</sup> G	+ 0.02 <sup>4</sup> F
21	$\Gamma_5$	31,558	0.60 <sup>2</sup> I	+	0.22 <sup>2</sup> H	+ 0.10 <sup>2</sup> F1
22	$\Gamma_{6,7}$	32,017	0.52 <sup>2</sup> I	+	0.18 <sup>2</sup> H	+ 0.14 <sup>4</sup> D
23	$\Gamma_{6,7}$	32,053	0.53 <sup>2</sup> I	+	0.31 <sup>2</sup> F1	+ 0.07 <sup>4</sup> G
24	$\Gamma_5$	32,341	0.90 <sup>4</sup> D	+	0.05 <sup>2</sup> I	+ 0.02 <sup>2</sup> F1
25	$\Gamma_{6,7}$	32,515	0.81 <sup>4</sup> D	+	0.12 <sup>2</sup> I	+ 0.04 <sup>2</sup> H
26	$\Gamma_5$	32,554	0.96 <sup>4</sup> D	+	0.02 <sup>2</sup> I	+ 0.01 <sup>2</sup> H
27	$\Gamma_{6,7}$	33,427	0.49 <sup>2</sup> I	+	0.20 <sup>2</sup> F1	+ 0.10 <sup>2</sup> H
28	$\Gamma_5$	33,591	0.48 <sup>2</sup> I	+	0.20 <sup>2</sup> F1	+ 0.10 <sup>4</sup> D
29	$\Gamma_{6,7}$	35,102	0.43 <sup>2</sup> I	+	0.25 <sup>2</sup> H	+ 0.12 <sup>2</sup> D3
30	$\Gamma_5$	38,794	0.39 <sup>2</sup> G2	+	0.20 <sup>2</sup> F1	+ 0.16 <sup>2</sup> F2
31	$\Gamma_{6,7}$	38,807	0.34 <sup>2</sup> G2	+	0.20 <sup>2</sup> F2	+ 0.16 <sup>2</sup> H
32	$\Gamma_5$	40,366	0.41 <sup>2</sup> I	+	0.15 <sup>4</sup> F	+ 0.15 <sup>4</sup> P
33	$\Gamma_5$	40,399	0.28 <sup>2</sup> F2	+	0.25 <sup>2</sup> G2	+ 0.11 <sup>2</sup> D2
34	$\Gamma_{6,7}$	40,492	0.29 <sup>2</sup> G2	+	0.21 <sup>2</sup> F2	+ 0.16 <sup>2</sup> I
35	$\Gamma_5$	41,114	0.30 <sup>4</sup> P	+	0.30 <sup>2</sup> I	+ 0.26 <sup>4</sup> F
36	$\Gamma_{6,7}$	41,115	0.39 <sup>4</sup> P	+	0.39 <sup>4</sup> F	+ 0.11 <sup>4</sup> G
37	$\Gamma_{6,7}$	41,246	0.40 <sup>4</sup> P	+	0.36 <sup>4</sup> F	+ 0.10 <sup>4</sup> G
38	$\Gamma_5$	41,368	0.38 <sup>4</sup> P	+	0.36 <sup>4</sup> F	+ 0.10 <sup>4</sup> G
39	$\Gamma_{6,7}$	43,563	0.42 <sup>2</sup> I	+	0.26 <sup>2</sup> D3	+ 0.10 <sup>2</sup> H
40	$\Gamma_5$	43,935	0.50 <sup>2</sup> G2	+	0.26 <sup>2</sup> I	+ 0.12 <sup>2</sup> G1
41	$\Gamma_{6,7}$	44,070	0.44 <sup>4</sup> F	+	0.28 <sup>2</sup> I	+ 0.19 <sup>2</sup> D3
42	$\Gamma_5$	44,497	0.43 <sup>2</sup> I	+	0.39 <sup>2</sup> D3	+ 0.06 <sup>2</sup> D1
43	$\Gamma_{6,7}$	44,820	0.55 <sup>4</sup> F	+	0.18 <sup>2</sup> D3	+ 0.15 <sup>2</sup> I
44	$\Gamma_{6,7}$	47,236	0.66 <sup>2</sup> G2	+	0.12 <sup>2</sup> D3	+ 0.05 <sup>4</sup> F
45	$\Gamma_5$	47,676	0.25 <sup>2</sup> F2	+	0.20 <sup>4</sup> F	+ 0.18 <sup>2</sup> G2
46	$\Gamma_{6,7}$	48,127	0.26 <sup>4</sup> F	+	0.23 <sup>2</sup> F2	+ 0.14 <sup>4</sup> G
47	$\Gamma_{6,7}$	48,789	0.48 <sup>4</sup> F	+	0.22 <sup>4</sup> G	+ 0.18 <sup>4</sup> P
48	$\Gamma_5$	49,092	0.52 <sup>4</sup> F	+	0.24 <sup>4</sup> G	+ 0.18 <sup>4</sup> P
49	$\Gamma_{6,7}$	49,185	0.30 <sup>2</sup> F2	+	0.23 <sup>2</sup> G2	+ 0.17 <sup>4</sup> F

Table 15. Energy levels ( $\text{cm}^{-1}$ ) for  $\text{Co}^{4+}$  using the parameters of tables 2 and 4, for Ge site in  $\text{Bi}_{12}\text{GeO}_{20}$ ,  $T$  symmetry,  $\zeta = 654 \text{ cm}^{-1}$  (cont'd)

No.	I.R. <sup>a</sup>	Energy (cm <sup>-1</sup> )	Free ion state				
50	Γ <sub>5</sub>	49,222	0.41 <sup>2</sup> F <sub>2</sub>	+	0.19 <sup>4</sup> F	+	0.12 <sup>2</sup> I
51	Γ <sub>6,7</sub>	49,356	0.43 <sup>2</sup> F <sub>2</sub>	+	0.29 <sup>2</sup> G <sub>2</sub>	+	0.09 <sup>4</sup> F
52	Γ <sub>5</sub>	49,398	0.49 <sup>2</sup> F <sub>2</sub>	+	0.14 <sup>4</sup> F	+	0.12 <sup>2</sup> I
53	Γ <sub>5</sub>	49,442	0.54 <sup>2</sup> F <sub>2</sub>	+	0.30 <sup>2</sup> G <sub>2</sub>	+	0.06 <sup>2</sup> I
54	Γ <sub>5</sub>	51,355	0.66 <sup>2</sup> H	+	0.24 <sup>2</sup> F <sub>1</sub>	+	0.03 <sup>4</sup> F
55	Γ <sub>6,7</sub>	51,366	0.63 <sup>2</sup> H	+	0.29 <sup>2</sup> F <sub>1</sub>	+	0.02 <sup>4</sup> F
56	Γ <sub>5</sub>	52,896	0.67 2S	+	0.24 <sup>2</sup> G <sub>2</sub>	+	0.04 <sup>2</sup> G <sub>1</sub>
57	Γ <sub>6,7</sub>	53,152	0.68 <sup>4</sup> F	+	0.17 <sup>4</sup> D	+	0.10 <sup>4</sup> G
58	Γ <sub>5</sub>	53,301	0.67 <sup>4</sup> F	+	0.16 <sup>4</sup> D	+	0.10 <sup>4</sup> G
59	Γ <sub>6,7</sub>	53,417	0.69 <sup>4</sup> F	+	0.19 <sup>4</sup> D	+	0.09 <sup>4</sup> G
60	Γ <sub>5</sub>	53,697	0.69 <sup>4</sup> F	+	0.20 <sup>4</sup> D	+	0.08 <sup>4</sup> G
61	Γ <sub>6,7</sub>	55,364	0.42 <sup>2</sup> H	+	0.30 <sup>2</sup> D <sub>3</sub>	+	0.22 <sup>2</sup> D <sub>2</sub>
62	Γ <sub>6,7</sub>	56,870	0.42 <sup>2</sup> H	+	0.18 <sup>2</sup> F <sub>1</sub>	+	0.16 <sup>2</sup> F <sub>2</sub>
63	Γ <sub>5</sub>	57,054	0.35 <sup>2</sup> H	+	0.24 <sup>2</sup> F <sub>1</sub>	+	0.14 <sup>2</sup> F <sub>2</sub>
64	Γ <sub>5</sub>	58,351	0.36 <sup>2</sup> F <sub>1</sub>	+	0.20 <sup>2</sup> H	+	0.14 <sup>2</sup> G <sub>2</sub>
65	Γ <sub>6,7</sub>	58,452	0.32 <sup>2</sup> F <sub>1</sub>	+	0.23 <sup>2</sup> H	+	0.12 <sup>2</sup> G <sub>2</sub>
66	Γ <sub>6,7</sub>	61,535	0.44 <sup>2</sup> H	+	0.26 <sup>2</sup> I	+	0.18 <sup>2</sup> F <sub>1</sub>
67	Γ <sub>5</sub>	61,781	0.56 <sup>2</sup> F <sub>1</sub>	+	0.22 <sup>2</sup> I	+	0.20 <sup>2</sup> F <sub>2</sub>
68	Γ <sub>5</sub>	61,964	0.46 <sup>2</sup> H	+	0.27 <sup>2</sup> I	+	0.16 <sup>2</sup> F <sub>1</sub>
69	Γ <sub>6,7</sub>	65,000	0.78 <sup>2</sup> G <sub>1</sub>	+	0.08 <sup>2</sup> H	+	0.07 <sup>2</sup> D <sub>1</sub>
70	Γ <sub>5</sub>	67,382	0.43 <sup>2</sup> D <sub>2</sub>	+	0.20 <sup>2</sup> G <sub>1</sub>	+	0.14 <sup>2</sup> I
71	Γ <sub>6,7</sub>	67,414	0.52 <sup>2</sup> D <sub>2</sub>	+	0.15 <sup>2</sup> G <sub>1</sub>	+	0.13 <sup>2</sup> I
72	Γ <sub>6,7</sub>	68,498	0.59 <sup>2</sup> D <sub>2</sub>	+	0.10 <sup>2</sup> H	+	0.10 <sup>2</sup> G <sub>2</sub>
73	Γ <sub>5</sub>	69,282	0.25 <sup>2</sup> D <sub>3</sub>	+	0.22 <sup>2</sup> H	+	0.18 <sup>2</sup> F <sub>1</sub>
74	Γ <sub>6,7</sub>	69,813	0.26 <sup>2</sup> D <sub>3</sub>	+	0.21 <sup>2</sup> H	+	0.17 <sup>2</sup> F <sub>1</sub>
75	Γ <sub>5</sub>	70,063	0.81 <sup>2</sup> G <sub>1</sub>	+	0.07 <sup>2</sup> P	+	0.05 <sup>2</sup> F <sub>2</sub>
76	Γ <sub>6,7</sub>	70,392	0.78 <sup>2</sup> G <sub>1</sub>	+	0.08 <sup>2</sup> P	+	0.06 <sup>2</sup> F <sub>2</sub>
77	Γ <sub>5</sub>	73,826	0.52 <sup>2</sup> G <sub>1</sub>	+	0.15 <sup>2</sup> F <sub>2</sub>	+	0.10 <sup>2</sup> G <sub>2</sub>
78	Γ <sub>6,7</sub>	73,964	0.51 <sup>2</sup> G <sub>1</sub>	+	0.13 <sup>2</sup> G <sub>2</sub>	+	0.11 <sup>2</sup> F <sub>2</sub>
79	Γ <sub>5</sub>	75,894	0.74 <sup>2</sup> G <sub>1</sub>	+	0.15 2S	+	0.10 <sup>2</sup> G <sub>2</sub>
80	Γ <sub>6,7</sub>	80,716	0.87 <sup>2</sup> P	+	0.05 <sup>2</sup> F <sub>2</sub>	+	0.04 <sup>2</sup> G <sub>2</sub>
81	Γ <sub>5</sub>	80,738	0.87 <sup>2</sup> P	+	0.04 <sup>2</sup> G <sub>2</sub>	+	0.04 <sup>2</sup> F <sub>2</sub>
82	Γ <sub>6,7</sub>	94,583	0.73 <sup>2</sup> D <sub>1</sub>	+	0.07 <sup>2</sup> D <sub>3</sub>	+	0.07 <sup>2</sup> G <sub>1</sub>
83	Γ <sub>5</sub>	94,621	0.72 <sup>2</sup> D <sub>1</sub>	+	0.07 <sup>2</sup> H	+	0.06 <sup>2</sup> D <sub>3</sub>
84	Γ <sub>6,7</sub>	94,712	0.73 <sup>2</sup> D <sub>1</sub>	+	0.08 <sup>2</sup> G <sub>1</sub>	+	0.08 <sup>2</sup> D <sub>3</sub>

<sup>a</sup>Irreducible representations of the double  $T$  (cubic) group (G. F. Koster, J. O. Dimmock, R. G. Wheeler, and H. Statz, *Properties of the thirty-two point groups*, MIT Press, Cambridge, MA (1963)).

**Table 16. Energy levels (cm<sup>-1</sup>) for Ni<sup>4+</sup> using the parameters of tables 2 and 4, for Ge site in Bi<sub>12</sub>GeO<sub>20</sub>, *T* symmetry, no spin orbit**

No.	I.R. <sup>a</sup>	Energy (cm <sup>-1</sup> )	Free ion state			
1	$\Gamma_{2,3}$	0	1.00 <sup>5</sup> D			
2	$\Gamma_4$	11,676	1.00 <sup>5</sup> D			
3	$\Gamma_4$	12,590	0.47 <sup>3</sup> H	+	0.17 <sup>3</sup> G	+ 0.13 <sup>3</sup> F2
4	$\Gamma_{2,3}$	21,226	0.87 <sup>3</sup> H	+	0.12 <sup>3</sup> G	+ 0.01 <sup>3</sup> D
5	$\Gamma_4$	22,495	0.42 <sup>3</sup> H	+	0.28 <sup>3</sup> G	+ 0.20 <sup>3</sup> P2
6	$\Gamma_4$	23,138	0.42 <sup>3</sup> F2	+	0.38 <sup>3</sup> H	+ 0.08 <sup>3</sup> F1
7	$\Gamma_4$	23,808	0.44 <sup>1</sup> I	+	0.27 <sup>1</sup> G2	+ 0.17 <sup>1</sup> G1
8	$\Gamma_1$	23,880	1.00 <sup>3</sup> G			
9	$\Gamma_{2,3}$	24,316	0.70 <sup>1</sup> I	+	0.16 <sup>1</sup> D2	+ 0.07 <sup>1</sup> G1
10	$\Gamma_1$	24,510	0.94 <sup>3</sup> F2	+	0.06 <sup>3</sup> F1	
11	$\Gamma_{2,3}$	26,499	0.49 <sup>3</sup> G	+	0.48 <sup>3</sup> D	+ 0.04 <sup>3</sup> H
12	$\Gamma_1$	31,376	0.32 <sup>1</sup> I	+	0.24 <sup>1</sup> G1	+ 0.23 <sup>1</sup> S2
13	$\Gamma_4$	31,437	0.55 <sup>3</sup> H	+	0.32 <sup>3</sup> G	+ 0.09 <sup>3</sup> P1
14	$\Gamma_1$	32,026	0.94 <sup>1</sup> I	+	0.06 <sup>1</sup> F	
15	$\Gamma_4$	32,626	0.57 <sup>1</sup> I	+	0.27 <sup>1</sup> G2	+ 0.07 <sup>1</sup> G1
16	$\Gamma_4$	33,069	0.37 <sup>3</sup> G	+	0.35 <sup>3</sup> H	+ 0.10 <sup>3</sup> F1
17	$\Gamma_4$	33,922	0.40 <sup>3</sup> F2	+	0.28 <sup>3</sup> H	+ 0.12 <sup>3</sup> G
18	$\Gamma_4$	34,628	0.44 <sup>1</sup> I	+	0.29 <sup>1</sup> G2	+ 0.14 <sup>1</sup> F
19	$\Gamma_4$	36,016	0.48 <sup>3</sup> D	+	0.23 <sup>3</sup> H	+ 0.15 <sup>3</sup> F2
20	$\Gamma_{2,3}$	36,226	0.47 <sup>1</sup> D2	+	0.34 <sup>1</sup> G2	+ 0.11 <sup>1</sup> I
21	$\Gamma_{2,3}$	36,905	0.52 <sup>3</sup> D	+	0.39 <sup>3</sup> G	+ 0.10 <sup>3</sup> H
22	$\Gamma_4$	37,571	0.44 <sup>3</sup> G	+	0.27 <sup>3</sup> D	+ 0.27 <sup>3</sup> F2
23	$\Gamma_4$	40,383	0.50 <sup>3</sup> P2	+	0.27 <sup>3</sup> F2	+ 0.14 <sup>3</sup> H
24	$\Gamma_4$	43,140	0.87 <sup>1</sup> I	+	0.11 <sup>1</sup> G2	+ 0.02 <sup>1</sup> G1
25	$\Gamma_1$	44,491	0.37 <sup>1</sup> I	+	0.32 <sup>1</sup> S2	+ 0.21 <sup>1</sup> G2
26	$\Gamma_4$	45,239	0.43 <sup>1</sup> F	+	0.30 <sup>1</sup> I	+ 0.18 <sup>1</sup> G2
27	$\Gamma_{2,3}$	45,314	0.48 <sup>1</sup> G2	+	0.18 <sup>1</sup> G1	+ 0.18 <sup>1</sup> I
28	$\Gamma_4$	46,520	0.69 <sup>1</sup> F	+	0.12 <sup>1</sup> D1	+ 0.08 <sup>1</sup> G1
29	$\Gamma_1$	48,325	0.57 <sup>1</sup> G2	+	0.20 <sup>1</sup> S2	+ 0.10 <sup>1</sup> G1
30	$\Gamma_1$	51,292	0.94 <sup>1</sup> F	+	0.06 <sup>1</sup> I	
31	$\Gamma_4$	51,301	0.73 <sup>1</sup> D2	+	0.09 <sup>1</sup> G1	+ 0.08 <sup>1</sup> F
32	$\Gamma_4$	51,464	0.51 <sup>1</sup> G2	+	0.42 <sup>1</sup> F	+ 0.06 <sup>1</sup> I
33	$\Gamma_4$	53,601	0.70 <sup>3</sup> F1	+	0.16 <sup>3</sup> P1	+ 0.10 <sup>3</sup> H
34	$\Gamma_1$	57,595	0.94 <sup>3</sup> F1	+	0.06 <sup>3</sup> F2	
35	$\Gamma_4$	58,214	0.53 <sup>3</sup> P1	+	0.17 <sup>3</sup> F2	+ 0.14 <sup>3</sup> P2
36	$\Gamma_4$	60,295	0.70 <sup>3</sup> F1	+	0.11 <sup>3</sup> D	+ 0.10 <sup>3</sup> G
37	$\Gamma_4$	63,745	0.77 <sup>1</sup> G1	+	0.20 <sup>1</sup> I	+ 0.02 <sup>1</sup> G2
38	$\Gamma_{2,3}$	63,749	0.74 <sup>1</sup> G1	+	0.17 <sup>1</sup> G2	+ 0.07 <sup>1</sup> D2
39	$\Gamma_1$	69,390	0.51 <sup>1</sup> G1	+	0.24 <sup>1</sup> I	+ 0.17 <sup>1</sup> S2
40	$\Gamma_4$	70,055	0.54 <sup>1</sup> G1	+	0.29 <sup>1</sup> G2	+ 0.10 <sup>1</sup> D2
41	$\Gamma_4$	76,978	0.75 <sup>1</sup> D1	+	0.11 <sup>1</sup> F	+ 0.06 <sup>1</sup> D2
42	$\Gamma_{2,3}$	79,827	0.84 <sup>1</sup> D1	+	0.16 <sup>1</sup> D2	
43	$\Gamma_1$	102,985	0.83 <sup>1</sup> S1	+	0.09 <sup>1</sup> S2	+ 0.08 <sup>1</sup> G1

<sup>a</sup>Irreducible representations of the *T* (cubic) group,  $\Gamma_{2,3} = \Gamma_2 + \Gamma_3$  (Bethe notation, G. F. Koster, J. O. Dimmock, R. G. Wheeler, and H. Statz, *Properties of the thirty-two point groups*, MIT Press, Cambridge, MA (1963)).



Table 17. Energy levels (cm<sup>-1</sup>) for Ni<sup>4+</sup> using the parameters of tables 2 and 4, for Ge site in Bi<sub>12</sub>GeO<sub>20</sub>, *T* symmetry,  $\zeta = 830$  cm<sup>-1</sup>

No.	I.R. <sup>a</sup>	Energy (cm <sup>-1</sup> )	Free ion state				
1	$\Gamma_1$	0	0.99 <sup>5</sup> D				
2	$\Gamma_4$	37	0.99 <sup>5</sup> D				
3	$\Gamma_{2,3}$	72	0.99 <sup>5</sup> D				
4	$\Gamma_4$	124	1.00 <sup>5</sup> D				
5	$\Gamma_1$	176	1.00 <sup>5</sup> D				
6	$\Gamma_4$	11,142	0.99 <sup>5</sup> D	+	0.01 <sup>3</sup> F2		
7	$\Gamma_4$	11,598	0.99 <sup>5</sup> D				
8	$\Gamma_{2,3}$	11,599	0.99 <sup>5</sup> D				
9	$\Gamma_4$	12,185	0.84 <sup>5</sup> D	+	0.06 <sup>3</sup> H	+	0.03 <sup>3</sup> G
10	$\Gamma_4$	12,230	0.99 <sup>5</sup> D				
11	$\Gamma_1$	12,247	0.99 <sup>5</sup> D				
12	$\Gamma_4$	12,400	0.38 <sup>3</sup> H	+	0.15 <sup>3</sup> G	+	0.15 <sup>5</sup> D
13	$\Gamma_{2,3}$	12,425	0.43 <sup>3</sup> H	+	0.19 <sup>3</sup> G	+	0.12 <sup>3</sup> F2
14	$\Gamma_4$	13,069	0.51 <sup>3</sup> H	+	0.14 <sup>3</sup> G	+	0.13 <sup>3</sup> F2
15	$\Gamma_1$	13,283	0.55 <sup>3</sup> H	+	0.13 <sup>3</sup> F2	+	0.12 <sup>3</sup> G
16	$\Gamma_4$	21,211	0.78 <sup>3</sup> H	+	0.11 <sup>3</sup> G	+	0.02 <sup>1</sup> G2
17	$\Gamma_4$	21,327	0.86 <sup>3</sup> H	+	0.13 <sup>3</sup> G	+	0.01 <sup>3</sup> D
18	$\Gamma_4$	21,985	0.41 <sup>3</sup> H	+	0.19 <sup>3</sup> G	+	0.16 <sup>3</sup> P2
19	$\Gamma_{2,3}$	22,258	0.47 <sup>3</sup> H	+	0.21 <sup>3</sup> G	+	0.19 <sup>3</sup> P2
20	$\Gamma_{2,3}$	22,738	0.29 <sup>3</sup> F2	+	0.27 <sup>3</sup> H	+	0.18 <sup>1</sup> I
21	$\Gamma_4$	22,802	0.36 <sup>3</sup> H	+	0.31 <sup>3</sup> F2	+	0.09 <sup>3</sup> G
22	$\Gamma_1$	22,857	0.40 <sup>3</sup> F2	+	0.37 <sup>3</sup> H	+	0.09 <sup>3</sup> F1
23	$\Gamma_4$	23,071	0.36 <sup>3</sup> H	+	0.34 <sup>3</sup> G	+	0.18 <sup>3</sup> P2
24	$\Gamma_1$	23,508	0.39 <sup>3</sup> G	+	0.31 <sup>3</sup> H	+	0.20 <sup>3</sup> P2
25	$\Gamma_4$	23,668	0.39 <sup>3</sup> F2	+	0.38 <sup>3</sup> H	+	0.07 <sup>3</sup> F1
26	$\Gamma_4$	23,960	0.97 <sup>3</sup> G	+	0.02 <sup>3</sup> H		
27	$\Gamma_4$	24,171	0.46 <sup>3</sup> F2	+	0.13 <sup>1</sup> I	+	0.12 <sup>3</sup> H
28	$\Gamma_4$	25,281	0.53 <sup>3</sup> F2	+	0.19 <sup>1</sup> I	+	0.08 <sup>1</sup> G2
29	$\Gamma_{2,3}$	25,307	0.50 <sup>1</sup> I	+	0.13 <sup>3</sup> F2	+	0.11 <sup>3</sup> H
30	$\Gamma_4$	26,669	0.48 <sup>3</sup> G	+	0.46 <sup>3</sup> D	+	0.05 <sup>3</sup> H
31	$\Gamma_4$	26,735	0.45 <sup>3</sup> D	+	0.44 <sup>3</sup> G	+	0.06 <sup>3</sup> H
32	$\Gamma_1$	30,260	0.25 <sup>3</sup> H	+	0.20 <sup>3</sup> G	+	0.13 <sup>1</sup> I
33	$\Gamma_4$	31,308	0.44 <sup>3</sup> H	+	0.22 <sup>3</sup> G	+	0.13 <sup>1</sup> I
34	$\Gamma_4$	31,620	0.55 <sup>3</sup> H	+	0.30 <sup>3</sup> G	+	0.10 <sup>3</sup> P1
35	$\Gamma_{2,3}$	31,780	0.56 <sup>3</sup> H	+	0.31 <sup>3</sup> G	+	0.07 <sup>3</sup> P1
36	$\Gamma_1$	31,787	0.68 <sup>1</sup> I	+	0.12 <sup>3</sup> H	+	0.08 <sup>3</sup> G
37	$\Gamma_4$	32,578	0.27 <sup>3</sup> H	+	0.26 <sup>3</sup> G	+	0.16 <sup>1</sup> I
38	$\Gamma_4$	32,789	0.41 <sup>3</sup> H	+	0.17 <sup>1</sup> I	+	0.11 <sup>1</sup> G1
39	$\Gamma_{2,3}$	33,217	0.34 <sup>3</sup> G	+	0.25 <sup>3</sup> H	+	0.20 <sup>3</sup> F2
40	$\Gamma_4$	33,309	0.29 <sup>3</sup> H	+	0.28 <sup>3</sup> G	+	0.12 <sup>3</sup> D
41	$\Gamma_1$	33,333	0.29 <sup>3</sup> H	+	0.26 <sup>3</sup> G	+	0.23 <sup>1</sup> I
42	$\Gamma_4$	33,701	0.25 <sup>3</sup> G	+	0.20 <sup>3</sup> H	+	0.17 <sup>1</sup> I
43	$\Gamma_{2,3}$	33,772	0.23 <sup>3</sup> H	+	0.21 <sup>3</sup> G	+	0.20 <sup>3</sup> F2
44	$\Gamma_4$	33,972	0.33 <sup>3</sup> F2	+	0.18 <sup>3</sup> H	+	0.15 <sup>3</sup> G
45	$\Gamma_4$	34,191	0.29 <sup>3</sup> H	+	0.28 <sup>3</sup> F2	+	0.11 <sup>3</sup> G
46	$\Gamma_1$	34,422	0.43 <sup>3</sup> F2	+	0.23 <sup>3</sup> H	+	0.09 <sup>3</sup> G
47	$\Gamma_4$	35,094	0.23 <sup>1</sup> I	+	0.15 <sup>1</sup> G2	+	0.12 <sup>3</sup> G
48	$\Gamma_{2,3}$	35,521	0.31 <sup>3</sup> D	+	0.22 <sup>3</sup> H	+	0.15 <sup>3</sup> F2
49	$\Gamma_4$	36,073	0.49 <sup>3</sup> D	+	0.26 <sup>3</sup> H	+	0.10 <sup>3</sup> F2

**Table 17. Energy levels (cm<sup>-1</sup>) for Ni<sup>4+</sup> using the parameters of tables 2 and 4, for Ge site in Bi<sub>12</sub>GeO<sub>20</sub>, *T* symmetry,  $\zeta = 830$  cm<sup>-1</sup> (cont'd)**

No.	I.R. <sup>a</sup>	Energy (cm <sup>-1</sup> )	Free ion state			
50	$\Gamma_4$	36,433	0.44 <sup>3</sup> D	+	0.21 <sup>3</sup> H	+ 0.17 <sup>3</sup> F2
51	$\Gamma_1$	36,794	0.41 <sup>3</sup> D	+	0.19 <sup>3</sup> H	+ 0.18 <sup>3</sup> F2
52	$\Gamma_4$	37,126	0.48 <sup>3</sup> D	+	0.41 <sup>3</sup> G	+ 0.10 <sup>3</sup> H
53	$\Gamma_{2,3}$	37,175	0.27 <sup>1</sup> D2	+	0.23 <sup>1</sup> G2	+ 0.13 <sup>3</sup> H
54	$\Gamma_4$	37,236	0.42 <sup>3</sup> D	+	0.39 <sup>3</sup> G	+ 0.08 <sup>3</sup> H
55	$\Gamma_{2,3}$	37,483	0.46 <sup>3</sup> G	+	0.25 <sup>3</sup> F2	+ 0.24 <sup>3</sup> D
56	$\Gamma_4$	37,610	0.41 <sup>3</sup> G	+	0.28 <sup>3</sup> F2	+ 0.22 <sup>3</sup> D
57	$\Gamma_4$	38,191	0.39 <sup>3</sup> G	+	0.32 <sup>3</sup> D	+ 0.26 <sup>3</sup> F2
58	$\Gamma_1$	38,510	0.37 <sup>3</sup> G	+	0.37 <sup>3</sup> D	+ 0.25 <sup>3</sup> F2
59	$\Gamma_4$	40,349	0.44 <sup>3</sup> P2	+	0.31 <sup>3</sup> F2	+ 0.15 <sup>3</sup> H
60	$\Gamma_{2,3}$	40,430	0.40 <sup>3</sup> P2	+	0.32 <sup>3</sup> F2	+ 0.14 <sup>3</sup> H
61	$\Gamma_4$	41,018	0.55 <sup>3</sup> P2	+	0.22 <sup>3</sup> F2	+ 0.12 <sup>3</sup> H
62	$\Gamma_1$	41,129	0.55 <sup>3</sup> P2	+	0.19 <sup>3</sup> F2	+ 0.12 <sup>3</sup> G
63	$\Gamma_4$	43,447	0.86 <sup>1</sup> I	+	0.11 <sup>1</sup> G2	+ 0.02 <sup>1</sup> G1
64	$\Gamma_1$	44,917	0.35 <sup>1</sup> I	+	0.30 <sup>1</sup> S2	+ 0.22 <sup>1</sup> G2
65	$\Gamma_4$	45,484	0.42 <sup>1</sup> F	+	0.30 <sup>1</sup> I	+ 0.18 <sup>1</sup> G2
66	$\Gamma_{2,3}$	45,638	0.48 <sup>1</sup> G2	+	0.17 <sup>1</sup> G1	+ 0.17 <sup>1</sup> I
67	$\Gamma_4$	46,727	0.67 <sup>1</sup> F	+	0.12 <sup>1</sup> D1	+ 0.07 <sup>1</sup> G1
68	$\Gamma_1$	48,344	0.54 <sup>1</sup> G2	+	0.19 <sup>1</sup> S2	+ 0.09 <sup>1</sup> G1
69	$\Gamma_4$	51,401	0.71 <sup>1</sup> D2	+	0.08 <sup>1</sup> G1	+ 0.08 <sup>1</sup> F
70	$\Gamma_1$	51,481	0.92 <sup>1</sup> F	+	0.06 <sup>1</sup> I	+ 0.01 <sup>3</sup> D
71	$\Gamma_4$	51,640	0.51 <sup>1</sup> G2	+	0.41 <sup>1</sup> F	+ 0.06 <sup>1</sup> I
72	$\Gamma_1$	53,599	0.63 <sup>3</sup> F1	+	0.19 <sup>3</sup> P1	+ 0.10 <sup>3</sup> H
73	$\Gamma_4$	53,659	0.67 <sup>3</sup> F1	+	0.19 <sup>3</sup> P1	+ 0.11 <sup>3</sup> H
74	$\Gamma_{2,3}$	54,208	0.72 <sup>3</sup> F1	+	0.13 <sup>3</sup> P1	+ 0.09 <sup>3</sup> H
75	$\Gamma_4$	54,282	0.72 <sup>3</sup> F1	+	0.12 <sup>3</sup> P1	+ 0.09 <sup>3</sup> H
76	$\Gamma_1$	57,747	0.50 <sup>3</sup> P1	+	0.19 <sup>3</sup> F2	+ 0.16 <sup>3</sup> F1
77	$\Gamma_4$	57,871	0.93 <sup>3</sup> F1	+	0.06 <sup>3</sup> F2	
78	$\Gamma_4$	58,101	0.51 <sup>3</sup> P1	+	0.18 <sup>3</sup> F2	+ 0.15 <sup>3</sup> F1
79	$\Gamma_{2,3}$	58,864	0.54 <sup>3</sup> P1	+	0.16 <sup>3</sup> P2	+ 0.15 <sup>3</sup> F2
80	$\Gamma_4$	59,063	0.53 <sup>3</sup> P1	+	0.15 <sup>3</sup> P2	+ 0.15 <sup>3</sup> F2
81	$\Gamma_{2,3}$	60,575	0.68 <sup>3</sup> F1	+	0.12 <sup>3</sup> D	+ 0.08 <sup>3</sup> G
82	$\Gamma_4$	60,614	0.70 <sup>3</sup> F1	+	0.11 <sup>3</sup> G	+ 0.10 <sup>3</sup> D
83	$\Gamma_1$	60,692	0.70 <sup>3</sup> F1	+	0.13 <sup>3</sup> G	+ 0.08 <sup>3</sup> D
84	$\Gamma_4$	60,754	0.69 <sup>3</sup> F1	+	0.12 <sup>3</sup> D	+ 0.08 <sup>3</sup> G
85	$\Gamma_4$	64,083	0.77 <sup>1</sup> G1	+	0.20 <sup>1</sup> I	+ 0.02 <sup>1</sup> G2
86	$\Gamma_{2,3}$	64,145	0.72 <sup>1</sup> G1	+	0.17 <sup>1</sup> G2	+ 0.07 <sup>1</sup> D2
87	$\Gamma_1$	69,754	0.51 <sup>1</sup> G1	+	0.24 <sup>1</sup> I	+ 0.17 <sup>1</sup> S2
88	$\Gamma_4$	70,343	0.53 <sup>1</sup> G1	+	0.29 <sup>1</sup> G2	+ 0.10 <sup>1</sup> D2
89	$\Gamma_4$	77,274	0.75 <sup>1</sup> D1	+	0.11 <sup>1</sup> F	+ 0.06 <sup>1</sup> D2
90	$\Gamma_{2,3}$	80,097	0.83 <sup>1</sup> D1	+	0.16 <sup>1</sup> D2	
91	$\Gamma_1$	103,295	0.83 <sup>1</sup> S1	+	0.09 <sup>1</sup> S2	+ 0.08 <sup>1</sup> G1

<sup>a</sup>Irreducible representations of the *T* group,  $\Gamma_{2,3} = \Gamma_2 + \Gamma_3$  (degenerate doublets) (G. F. Koster, J. O. Dimmock, R. G. Wheeler, and H. Statz, *Properties of the thirty-two point groups*, MIT Press, Cambridge, MA (1963)).

**Table 18. Energy levels (cm<sup>-1</sup>) for Cu<sup>4+</sup> using the parameters of tables 2 and 4, for Ge site in Bi<sub>12</sub>GeO<sub>20</sub>, *T* symmetry, no spin orbit**

No.	I.R. <sup>a</sup>	Energy (cm <sup>-1</sup> )	Free ion state
1	$\Gamma_1$	0	1.00 <sup>4</sup> F
2	$\Gamma_4$	9,490	1.00 <sup>4</sup> F
3	$\Gamma_4$	14,930	0.76 <sup>4</sup> F + 0.24 <sup>4</sup> P
4	$\Gamma_{2,3}$	18,807	0.66 <sup>2</sup> G + 0.19 <sup>2</sup> H + 0.14 <sup>2</sup> D <sub>2</sub>
5	$\Gamma_4$	19,357	0.46 <sup>2</sup> G + 0.34 <sup>2</sup> H + 0.18 <sup>2</sup> P
6	$\Gamma_4$	23,777	0.76 <sup>4</sup> P + 0.24 <sup>4</sup> F
7	$\Gamma_4$	25,491	0.31 <sup>2</sup> H + 0.29 <sup>2</sup> G + 0.23 <sup>2</sup> D <sub>2</sub>
8	$\Gamma_1$	26,206	1.00 <sup>2</sup> G
9	$\Gamma_4$	28,207	0.51 <sup>2</sup> G + 0.48 <sup>2</sup> H
10	$\Gamma_4$	28,786	0.53 <sup>2</sup> H + 0.24 <sup>2</sup> G + 0.23 <sup>2</sup> P
11	$\Gamma_{2,3}$	30,961	0.49 <sup>2</sup> H + 0.44 <sup>2</sup> D <sub>2</sub> + 0.07 <sup>2</sup> D <sub>1</sub>
12	$\Gamma_4$	32,785	0.43 <sup>2</sup> P + 0.38 <sup>2</sup> H + 0.17 <sup>2</sup> F
13	$\Gamma_4$	37,536	0.55 <sup>2</sup> D <sub>2</sub> + 0.26 <sup>2</sup> F + 0.09 <sup>2</sup> G
14	$\Gamma_4$	37,953	0.72 <sup>2</sup> H + 0.28 <sup>2</sup> G
15	$\Gamma_1$	39,854	1.00 <sup>2</sup> F
16	$\Gamma_4$	41,737	0.66 <sup>2</sup> F + 0.12 <sup>2</sup> G + 0.10 <sup>2</sup> H
17	$\Gamma_{2,3}$	43,525	0.40 <sup>2</sup> D <sub>2</sub> + 0.32 <sup>2</sup> G + 0.28 <sup>2</sup> H
18	$\Gamma_4$	45,523	0.81 <sup>2</sup> F + 0.16 <sup>2</sup> P + 0.03 <sup>2</sup> H
19	$\Gamma_{2,3}$	64,919	0.91 <sup>2</sup> D <sub>1</sub> + 0.04 <sup>2</sup> H + 0.03 <sup>2</sup> G
20	$\Gamma_4$	65,096	0.76 <sup>2</sup> D <sub>1</sub> + 0.16 <sup>2</sup> D <sub>2</sub> + 0.05 <sup>2</sup> F

<sup>a</sup>Irreducible representations of the *T* group,  $\Gamma_{2,3} = \Gamma_2 + \Gamma_3$  (Bethe notation, G. F. Koster, J. O. Dimmock, R. G. Wheeler, and H. Statz, *Properties of the thirty-two point groups*, MIT Press, Cambridge, MA (1963)).

Table 19. Energy levels (cm<sup>-1</sup>) for Cu<sup>4+</sup> using the parameters of tables 2 and 4, for Ge site in Bi<sub>12</sub>GeO<sub>20</sub>, *T* symmetry,  $\zeta = 1008$  cm<sup>-1</sup>

No.	I.R. <sup>a</sup>	Energy (cm <sup>-1</sup> )	Free ion state			
1	$\Gamma_{6,7}$	0	1.00 <sup>4</sup> F			
2	$\Gamma_{6,7}$	9,024	0.98 <sup>4</sup> F	+	0.01 <sup>2</sup> G	
3	$\Gamma_5$	9,345	0.99 <sup>4</sup> F	+	0.01 <sup>2</sup> G	
4	$\Gamma_{6,7}$	9,818	0.99 <sup>4</sup> F			
5	$\Gamma_5$	10,072	1.00 <sup>4</sup> F			
6	$\Gamma_5$	14,203	0.81 <sup>4</sup> F	+	0.16 <sup>4</sup> P	+ 0.02 <sup>2</sup> G
7	$\Gamma_{6,7}$	14,782	0.75 <sup>4</sup> F	+	0.21 <sup>4</sup> P	+ 0.02 <sup>2</sup> G
8	$\Gamma_5$	15,460	0.69 <sup>4</sup> F	+	0.30 <sup>4</sup> P	+ 0.01 <sup>2</sup> D <sub>2</sub>
9	$\Gamma_{6,7}$	15,686	0.69 <sup>4</sup> F	+	0.30 <sup>4</sup> P	
10	$\Gamma_{6,7}$	18,892	0.62 <sup>2</sup> G	+	0.22 <sup>2</sup> H	+ 0.11 <sup>2</sup> D <sub>2</sub>
11	$\Gamma_5$	19,255	0.46 <sup>2</sup> G	+	0.35 <sup>2</sup> H	+ 0.14 <sup>2</sup> P
12	$\Gamma_{6,7}$	19,813	0.40 <sup>2</sup> G	+	0.29 <sup>2</sup> H	+ 0.20 <sup>2</sup> P
13	$\Gamma_{6,7}$	23,746	0.69 <sup>4</sup> P	+	0.28 <sup>4</sup> F	+ 0.01 <sup>2</sup> D <sub>2</sub>
14	$\Gamma_{6,7}$	23,816	0.73 <sup>4</sup> P	+	0.20 <sup>4</sup> F	+ 0.04 <sup>2</sup> G
15	$\Gamma_5$	23,834	0.69 <sup>4</sup> P	+	0.30 <sup>4</sup> F	
16	$\Gamma_5$	24,145	0.79 <sup>4</sup> P	+	0.14 <sup>4</sup> F	+ 0.05 <sup>2</sup> G
17	$\Gamma_5$	25,520	0.41 <sup>2</sup> H	+	0.22 <sup>2</sup> D <sub>2</sub>	+ 0.19 <sup>2</sup> G
18	$\Gamma_{6,7}$	25,870	0.40 <sup>2</sup> G	+	0.23 <sup>2</sup> H	+ 0.21 <sup>2</sup> D <sub>2</sub>
19	$\Gamma_5$	26,782	0.93 <sup>2</sup> G	+	0.03 <sup>4</sup> F	+ 0.02 <sup>4</sup> P
20	$\Gamma_{6,7}$	28,294	0.59 <sup>2</sup> H	+	0.38 <sup>2</sup> G	+ 0.01 <sup>2</sup> D <sub>1</sub>
21	$\Gamma_5$	28,589	0.57 <sup>2</sup> G	+	0.40 <sup>2</sup> H	+ 0.02 <sup>2</sup> D <sub>2</sub>
22	$\Gamma_{6,7}$	29,042	0.55 <sup>2</sup> H	+	0.25 <sup>2</sup> G	+ 0.16 <sup>2</sup> P
23	$\Gamma_5$	29,773	0.54 <sup>2</sup> H	+	0.25 <sup>2</sup> P	+ 0.18 <sup>2</sup> G
24	$\Gamma_{6,7}$	31,342	0.48 <sup>2</sup> H	+	0.40 <sup>2</sup> D <sub>2</sub>	+ 0.06 <sup>2</sup> D <sub>1</sub>
25	$\Gamma_{6,7}$	33,225	0.40 <sup>2</sup> P	+	0.34 <sup>2</sup> H	+ 0.14 <sup>2</sup> F
26	$\Gamma_5$	33,338	0.40 <sup>2</sup> P	+	0.38 <sup>2</sup> H	+ 0.18 <sup>2</sup> F
27	$\Gamma_5$	37,543	0.52 <sup>2</sup> D <sub>2</sub>	+	0.24 <sup>2</sup> F	+ 0.10 <sup>2</sup> G
28	$\Gamma_{6,7}$	37,782	0.40 <sup>2</sup> D <sub>2</sub>	+	0.25 <sup>2</sup> H	+ 0.22 <sup>2</sup> F
29	$\Gamma_{6,7}$	38,508	0.54 <sup>2</sup> H	+	0.23 <sup>2</sup> G	+ 0.14 <sup>2</sup> D <sub>2</sub>
30	$\Gamma_5$	38,664	0.68 <sup>2</sup> H	+	0.31 <sup>2</sup> G	+ 0.01 <sup>2</sup> P
31	$\Gamma_5$	40,084	1.00 <sup>2</sup> F			
32	$\Gamma_{6,7}$	41,764	0.61 <sup>2</sup> F	+	0.11 <sup>2</sup> H	+ 0.11 <sup>2</sup> G
33	$\Gamma_5$	42,706	0.68 <sup>2</sup> F	+	0.14 <sup>2</sup> G	+ 0.07 <sup>2</sup> H
34	$\Gamma_{6,7}$	43,857	0.37 <sup>2</sup> D <sub>2</sub>	+	0.30 <sup>2</sup> G	+ 0.27 <sup>2</sup> H
35	$\Gamma_{6,7}$	45,756	0.76 <sup>2</sup> F	+	0.16 <sup>2</sup> P	+ 0.04 <sup>2</sup> H
36	$\Gamma_5$	46,137	0.79 <sup>2</sup> F	+	0.17 <sup>2</sup> P	+ 0.03 <sup>2</sup> H
37	$\Gamma_5$	65,145	0.76 <sup>2</sup> D <sub>1</sub>	+	0.18 <sup>2</sup> D <sub>2</sub>	+ 0.04 <sup>2</sup> F
38	$\Gamma_{6,7}$	65,303	0.90 <sup>2</sup> D <sub>1</sub>	+	0.04 <sup>2</sup> H	+ 0.03 <sup>2</sup> D <sub>2</sub>
39	$\Gamma_{6,7}$	65,692	0.77 <sup>2</sup> D <sub>1</sub>	+	0.14 <sup>2</sup> D <sub>2</sub>	+ 0.06 <sup>2</sup> F

<sup>a</sup>Irreducible representations of the double *T* (cubic) group,  $\Gamma_{6,7} = \Gamma_6 + \Gamma_7$  (G. F. Koster, J. O. Dimmock, R. G. Wheeler, and H. Statz, *Properties of the thirty-two point groups*, MIT Press, Cambridge, MA (1963)).

**Table 20. Energy levels (cm<sup>-1</sup>) for Zn<sup>4+</sup> using the parameters of tables 2 and 4, for Ge site in Bi<sub>12</sub>GeO<sub>20</sub>, *T* symmetry, no spin orbit**

No.	I.R. <sup>a</sup>	Energy (cm <sup>-1</sup> )	Free ion state		
1	$\Gamma_4$	0	0.96 <sup>3</sup> F	+	0.04 <sup>3</sup> P
2	$\Gamma_4$	6,548	1.00 <sup>3</sup> F		
3	$\Gamma_1$	14,021	1.00 <sup>3</sup> F		
4	$\Gamma_4$	14,327	0.73 <sup>1</sup> D	+	0.27 <sup>1</sup> G
5	$\Gamma_{2,3}$	14,713	0.59 <sup>1</sup> D	+	0.41 <sup>1</sup> G
6	$\Gamma_4$	16,258	0.96 <sup>3</sup> P	+	0.04 <sup>3</sup> F
7	$\Gamma_4$	22,686	0.73 <sup>1</sup> G	+	0.27 <sup>1</sup> D
8	$\Gamma_1$	24,435	0.94 <sup>1</sup> G	+	0.06 <sup>1</sup> S
9	$\Gamma_4$	24,725	1.00 <sup>1</sup> G		
10	$\Gamma_{2,3}$	29,774	0.59 <sup>1</sup> G	+	0.41 <sup>1</sup> D
11	$\Gamma_1$	56,276	0.94 <sup>1</sup> S	+	0.06 <sup>1</sup> G

**Table 21. Energy levels (cm<sup>-1</sup>) for Zn<sup>4+</sup> using the parameters of tables 2 and 4, for Ge site in Bi<sub>12</sub>GeO<sub>20</sub>, *T* symmetry,  $\zeta = 1203$  cm<sup>-1</sup>**

No.	I.R. <sup>a</sup>	Energy (cm <sup>-1</sup> )	Free ion state				
1	$\Gamma_1$	0	0.97 <sup>3</sup> F	+	0.03 <sup>3</sup> P		
2	$\Gamma_4$	699	0.97 <sup>3</sup> F	+	0.03 <sup>3</sup> P		
3	$\Gamma_{2,3}$	1,877	0.96 <sup>3</sup> F	+	0.03 <sup>3</sup> P		
4	$\Gamma_4$	2,325	0.95 <sup>3</sup> F	+	0.04 <sup>3</sup> P		
5	$\Gamma_4$	8,121	0.95 <sup>3</sup> F	+	0.03 <sup>1</sup> D	+	0.01 <sup>1</sup> G
6	$\Gamma_4$	8,182	1.00 <sup>3</sup> F				
7	$\Gamma_{2,3}$	8,525	0.98 <sup>3</sup> F	+	0.01 <sup>1</sup> D	+	0.01 <sup>3</sup> P
8	$\Gamma_1$	8,926	1.00 <sup>3</sup> F				
9	$\Gamma_4$	15,124	0.40 <sup>1</sup> D	+	0.39 <sup>3</sup> F	+	0.15 <sup>3</sup> P
10	$\Gamma_{2,3}$	15,327	0.39 <sup>1</sup> D	+	0.39 <sup>3</sup> P	+	0.22 <sup>1</sup> G
11	$\Gamma_4$	16,486	0.65 <sup>3</sup> F	+	0.17 <sup>3</sup> P	+	0.10 <sup>1</sup> D
12	$\Gamma_4$	18,312	0.62 <sup>3</sup> P	+	0.19 <sup>1</sup> D	+	0.16 <sup>1</sup> G
13	$\Gamma_4$	18,589	0.97 <sup>3</sup> P	+	0.03 <sup>3</sup> F		
14	$\Gamma_{2,3}$	18,701	0.57 <sup>3</sup> P	+	0.20 <sup>1</sup> G	+	0.18 <sup>1</sup> D
15	$\Gamma_1$	18,903	0.96 <sup>3</sup> P	+	0.03 <sup>3</sup> F	+	0.01 <sup>1</sup> S
16	$\Gamma_4$	24,734	0.69 <sup>1</sup> G	+	0.28 <sup>1</sup> D	+	0.02 <sup>3</sup> F
17	$\Gamma_1$	26,307	0.94 <sup>1</sup> G	+	0.05 <sup>1</sup> S	+	0.01 <sup>3</sup> P
18	$\Gamma_4$	26,565	1.00 <sup>1</sup> G				
19	$\Gamma_{2,3}$	31,744	0.58 <sup>1</sup> G	+	0.41 <sup>1</sup> D	+	0.01 <sup>3</sup> P
20	$\Gamma_1$	58,264	0.94 <sup>1</sup> S	+	0.06 <sup>1</sup> G	+	0.01 <sup>3</sup> P

<sup>a</sup>Irreducible representations of the *T* group,  $\Gamma_{2,3} = \Gamma_2 + \Gamma_3$  (degenerate doublets) (G. F. Koster, J. O. Dimmock, R. G. Wheeler, and H. Statz, *Properties of the thirty-two point groups*, MIT Press, Cambridge, MA (1963)).

**Table 22. Energy levels (cm<sup>-1</sup>) for Ga<sup>4+</sup> using the parameters of tables 2 and 4, for Ge site in Bi<sub>12</sub>GeO<sub>20</sub>, *T* symmetry, no spin orbit**

No.	I.R. <sup>a</sup>	Energy (cm <sup>-1</sup> )	Free ion state
1	$\Gamma_4$	0	1.00 <sup>2D</sup>
2	$\Gamma_{2,3}$	4,981	1.00 <sup>2D</sup>

<sup>a</sup>Irreducible representations of the *T* (cubic) group,  $\Gamma_{2,3} = \Gamma_2 + \Gamma_3$  (Bethe notation, G. F. Koster, J. O. Dimmock, R. G. Wheeler, and H. Statz, *Properties of the thirty-two point groups*, MIT Press, Cambridge, MA (1963)).

**Table 23. Energy levels (cm<sup>-1</sup>) for Ga<sup>4+</sup> using the parameters of tables 2 and 4, for Ge site in Bi<sub>12</sub>GeO<sub>20</sub>, *T* symmetry,  $\zeta = 1496$  cm<sup>-1</sup>**

No.	I.R. <sup>a</sup>	Energy (cm <sup>-1</sup> )	Free ion state
1	$\Gamma_5$	0	1.00 <sup>2D</sup>
2	$\Gamma_{6,7}$	1,561	1.00 <sup>2D</sup>
3	$\Gamma_{6,7}$	7,160	1.00 <sup>2D</sup>

<sup>a</sup>Irreducible representations of the double *T* (cubic) group,  $\Gamma_{6,7} = \Gamma_6 + \Gamma_7$  (G. F. Koster, J. O. Dimmock, R. G. Wheeler, and H. Statz, *Properties of the thirty-two point groups*, MIT Press, Cambridge, MA (1963)).

## 6. Conclusion

The crystallographic data on six host materials have been used to find the crystal-field components at sites that could be occupied by quadruply ionized transition-metal ions with the  $3d^N$  electronic configuration. Using approximate values for the effective radial integrals  $\rho_2$  and  $\rho_4$ , we obtain the crystal-field parameters,  $B_{kq}$ , for Cr<sup>4+</sup> in the host crystals NaTiSiO<sub>5</sub>, Y<sub>2</sub>SiBeO<sub>7</sub>, Bi<sub>4</sub>Si<sub>3</sub>O<sub>12</sub>, and Bi<sub>4</sub>Ge<sub>3</sub>O<sub>12</sub>, assuming that the Cr<sup>4+</sup> ion occupied the Ge, Ti, or Si site in these crystals.

For the host crystal Bi<sub>12</sub>GeO<sub>20</sub> the Ge site has the cubic group symmetry *T*, and the entire quadruply ionized  $3d^N$  series energy levels were calculated using estimated crystal-field parameters. The calculated energy levels are for  $\zeta = 0$  and for the value of  $\zeta$  corresponding to the free-ion results. Experimental values of several energy levels for Cr<sup>4+</sup> have been reported and upon comparison with the results obtained here, our estimated crystal-field parameters are too large. More experimental data are needed to ascertain whether this difference is significant.

## 7. References

- (1) D. T. Sviridov, R. K. Sviridova, N. I. Kulik, and V. B. Glasko, *Optical spectra of the iso-electronic ions  $V^{2+}$ ,  $Cr^{3+}$ , and  $Mn^{4+}$  in an octahedral coordination*, J. Appl. Spectrosc. **30** (1979), 334.
- (2) S. Fraga, K. M. S. Saxena, and J. Karwowski, *Physical science data: 5. Handbook of atomic data*, Elsevier, New York (1976).
- (3) P. H. M. Uylings, A. J. J. Raassen, and J. F. Wyart, *Energies of  $N$  equivalent electrons expressed in terms of two-electron energies and independent three-electron parameters: A new complete set of orthogonal operators: II. Application of  $3d^N$  configurations*, J. Phys. **B17** (1984).
- (4) C. A. Morrison, *Host materials for transition-metal ions*, Harry Diamond Laboratories, HDL-DS-89-1 (September 1989). Pages referenced in the present report are to specific items. This data summary (HDL-DS-89-1) includes results based on the work of many researchers, and references to the researcher's work are given within it.
- (5) C. W. Nielson and G. F. Koster, *Spectroscopic coefficients for the  $p^n$ ,  $d^n$ , and  $f^n$  configurations*, MIT Press, Cambridge, MA (1963).
- (6) E. Konig and S. Kremer, *Ligand field energy diagrams*, Plenum Press, New York, NY (1977).
- (7) C. J. Ballhausen, *Ligand field theory*, McGraw Hill, New York (1962), p 93.
- (8) J. S. Griffith, *The theory of transition-metal ions*, Cambridge University Press, Cambridge, England (1961), p 226.
- (9) M. Weber, *CRC Handbook of laser science and technology*, Vol. III, p 198; Vol. IV, p 370; Vol. V, pp 12, 75, 78–80, 172, CRC Press, Boca Raton, FL (1987).
- (10) W. Wardzynski, T. Lukasiewicz, and J. Zmija, *Reversible photochromic effects in doped single crystals of bismuth germinate ( $Bi_{12}GeO_{20}$ ) and bismuth silicon oxide ( $Bi_{12}SiO_{20}$ )*, Opt. Commun. **30** (1979), 203.
- (11) W. Wardzynski, H. Szymczak, K. Pataj, T. Lukasiewicz, and J. Zmija, *Light induced charge transfer in Cr doped  $Bi_{12}GeO_{20}$  and  $Bi_{12}SiO_{20}$  single crystals*, J. Phys. Chem. Solids **43** (1982), 767.
- (12) G. F. Koster, J. O. Dimmock, R. G. Wheeler, and H. Statz, *Properties of the thirty-two point groups*, MIT Press, Cambridge, MA (1963).

- (13) W. Wardzynski, M. Baran, and H. Szymczak, *Electron paramagnetic resonance of  $Fe^{3+}$  in bismuth germanium oxide single crystals*, Physica **111B** (1981), 47.
- (14) W. Wardzynski and H. Szymczak, *The center of orthorhombic symmetry in chromium doped  $Bi_{12}GeO_{20}$  and  $Bi_{12}SiO_{20}$  single crystals*, J. Phys. Chem. Solids **45** (1984), 887.



## **Appendix A.—Crystallographic and X-Ray Data**

## Contents

	Page
A-1. $\text{Na}_2\text{TiSiO}_5$ .....	43
A-1.1 Crystallographic Data on $\text{Na}_2\text{TiSiO}_5$ .....	43
A-1.2 Crystal-Field Components, $A_{kq}$ , for the Si ( $D_{2d}$ ) Site in $\text{Na}_2\text{TiSiO}_5$ .....	43
A-1.3 Crystal-Field Components, $A_{kq}$ , for the Ti ( $C_{4v}$ ) Site in $\text{Na}_2\text{TiSiO}_5$ .....	43
A-1.4 $\text{Na}_2\text{TiSiO}_5$ References.....	44
A-2. $\text{Y}_2\text{SiBe}_2\text{O}_7$ .....	44
A-2.1 Crystallographic Data on $\text{Y}_2\text{SiBe}_2\text{O}_7$ .....	44
A-2.2 Crystal-Field Components, $A_{kq}$ , for the Si Site ( $S_4$ ) in $\text{Y}_2\text{SiBe}_2\text{O}_7$ .....	44
A-2.3 $\text{Y}_2\text{SiBe}_2\text{O}_7$ References.....	44
A-3. $\text{Bi}_4\text{X}_3\text{O}_{12}$ ( $X = \text{Si}, \text{Ge}$ ).....	45
A-3.1 Crystallographic Data on $\text{Bi}_4\text{X}_3\text{O}_{12}$ .....	45
A-3.2 X-Ray Data on $\text{Bi}_4\text{X}_3\text{O}_{12}$ .....	45
A-3.3 Crystal-Field Components, $A_{kq}$ , for the Si Site ( $S_4$ ) of $\text{Bi}_4\text{Si}_3\text{O}_{12}$ .....	45
A-3.4 Crystal-Field Components, $A_{kq}$ for the Ge Site ( $S_4$ ) of $\text{Bi}_4\text{Ge}_3\text{O}_{12}$ .....	46
A-3.5 $\text{Bi}_4\text{X}_3\text{O}_{12}$ References.....	46
A-4. $\text{Bi}_{12}\text{XO}_{20}$ ( $X = \text{Ge}, \text{Si}$ ).....	47
A-4.1 Crystallographic Data on $\text{Bi}_{12}\text{XO}_{20}$ .....	47
A-4.2 X-Ray Data on $\text{Bi}_{12}\text{XO}_{20}$ .....	48
A-4.3 Crystal-Field Components, $A_{kq}$ , for the Ge Site of $\text{Bi}_{12}\text{GeO}_{20}$ .....	48
A-4.4 Crystal-Field Components, $A_{kq}$ , for the Si Site ( $T$ ) of $\text{Bi}_{12}\text{SiO}_{20}$ .....	48
A-4.5 $\text{Bi}_{12}\text{XO}_{20}$ ( $X = \text{Ge}, \text{Si}$ ) References.....	48

The following tables give the crystallographic and x-ray data on each of the compounds considered. The crystal-field components,  $A_{kq}$ , are calculated for the monopole (point-charge), self-induced, and dipole contributions. Following each table is a list of references for that particular compound. These references include such topics as crystal growth, index of refraction, and electron spin resonance investigations. However, the list of references is not, in any sense, exhaustive and if further investigations are contemplated on any compound, additional library searches should be undertaken.

## A-1. $\text{Na}_2\text{TiSiO}_5$

### A-1.1 Crystallographic Data on $\text{Na}_2\text{TiSiO}_5$

Tetragonal ( $P4/nmm$ ), 129,  $Z = 2^a$

Ion	Site	Symmetry	$x^b$	$y$	$z$	$q$	$\alpha$ ( $\text{\AA}^3$ ) <sup>c</sup>
Ti	2c	$C_{4v}$	1/2	0	0.9343	4	0.506
Na	4e	$C_{2h}$	1/4	1/4	1/2	1	0.147
Si	2a	$D_{2d}$	0	0	0	4	0.0165
O <sub>1</sub>	8i	$C_s$	0	0.2071	0.1831	-2	1.349
O <sub>2</sub>	2c	$C_{4v}$	1/2	0	-0.7338	-2	1.349

<sup>a</sup>N. F. M. Henry and K. Lonsdale, *International tables for x-ray crystallography*, Vol. I: Symmetry groups, Kynoch, Birmingham, UK (1969).

<sup>b</sup>X-ray data:  $a = 6.480$   $\text{\AA}$ ,  $c = 5.107$   $\text{\AA}$  (Nyman et al (1978)).

<sup>c</sup>Schmidt et al (1979).

### A-1.2 Crystal-Field Components, $A_{kq}$ ( $\text{cm}^{-1}/\text{\AA}^k$ ) for the Si ( $D_{2d}$ ) Site in $\text{Na}_2\text{TiSiO}_5$

$A_{kq}$	Monopole	Self-induced	Dipole	Total
$A_{20}$	7,285	548.7	-34,349	-26,515
$A_{32}$	-64,975	25,258	-48,610	-88,327
$A_{40}$	-31,427	19,391	-20,545	-32,581
$A_{44}$	17,004	-12,063	21,554	26,494
$A_{52}$	326.3	-628.4	9,042	8,740

### A-1.3 Crystal-Field Components, $A_{kq}$ ( $\text{cm}^{-1}/\text{\AA}^k$ ), for the Ti ( $C_{4v}$ ) Site in $\text{Na}_2\text{TiSiO}_5$

$A_{kq}$	Monopole	Self-induced	Dipole	Total
$A_{10}$	15,041	0	114,675	129,716
$A_{20}$	18,952	-2,906	38,432	54,478
$A_{30}$	48,794	-13,854	36,992	71,932
$A_{40}$	18,057	-10,303	42,160	49,914
$A_{44}$	10,890	-4,552	766.4	7,105
$A_{50}$	4,284	-5,673	25,232	23,843
$A_{54}$	-6,132	3,088	2,933	-111.3

### A-1.4 Na<sub>2</sub>TiSiO<sub>5</sub> References

Nyman, H., M. O'Keeffe, and J. O. Borin, *Sodium titanium silicates, Na<sub>2</sub>TiSiO<sub>5</sub>*, Acta. Crystallogr. B34 (1978), 905.

Schmidt, P. C, A. Weiss, and T. P. Das, *Effect of crystal-fields and self-consistency on dipole and quadrupole polarizabilities of closed shell ions*, Phys. Rev. B19 (1979), 5525.

## A-2. Y<sub>2</sub>SiBe<sub>2</sub>O<sub>7</sub>

### A-2.1 Crystallographic Data on Y<sub>2</sub>SiBe<sub>2</sub>O<sub>7</sub>

Tetragonal  $D_{3d}^3$  ( $P\bar{4}2_1m$ ), 113,  $Z = 2$

Ion	Site	Symmetry	$x^a$	$y$	$z$	$q$	$\alpha$ (Å <sup>3</sup> ) <sup>b</sup>
Y	4e	C <sub>2</sub>	0.1595	0.6595	0.4873	3	0.870
Si	2a	S <sub>4</sub>	0	0	0	4	0.0165 <sup>c</sup>
Be	4e	C <sub>2</sub>	0.363	0.863	0.031	2	0.0125
O <sub>1</sub>	8f	C <sub>1</sub>	0.0823	0.1664	0.7928	-2	1.349
O <sub>2</sub>	4e	C <sub>2</sub>	0.3561	0.8561	0.7053	-2	1.349
O <sub>3</sub>	2c	C <sub>2v</sub>	0	1/2	0.8275	-2	1.349

<sup>a</sup>X-ray data:  $a = 7.283$  Å;  $c = 4.755$  Å, Bartram (1969).

<sup>b</sup>Schmidt et al (1979).

<sup>c</sup>Tessman et al (1953).

### A-2.2 Crystal-Field Components, $A_{kq}$ (cm<sup>-1</sup>/Å<sup>k</sup>), for the Si Site (S<sub>4</sub>) in Y<sub>2</sub>SiBe<sub>2</sub>O<sub>7</sub>

$A_{kq}$	Monopole	Self-induced	Dipole	Total
A <sub>20</sub>	10,029	-468.9	-19,678	-10,118
ReA <sub>32</sub>	35,565	-12,974	26,656	49,247
ImA <sub>32</sub>	-46,373	17,029	-40,428	-69,772
A <sub>40</sub>	-28,792	16,971	-23,527	-35,348
ReA <sub>44</sub>	-4,073	2,468	-6,484	-8,089
ImA <sub>44</sub>	-14,462	9,193	-17,375	-22,644
ReA <sub>52</sub>	902.0	-662.1	-2,652	-2,412
ImA <sub>52</sub>	-1,164	866.0	3,394	3,096
A <sub>44</sub>	15,025	—	—	24,045

### A-2.3 Y<sub>2</sub>SiBe<sub>2</sub>O<sub>7</sub> References

Bartram, S. F., *Crystal structure of Y<sub>2</sub>SiBe<sub>2</sub>O<sub>7</sub>*, Acta. Crystallogr. B25 (1969), 791.

Morrison, C. A., N. Karayianis, and D. E. Wortman, *Rare-earth ion-host interactions: 10. Lanthanides in Y<sub>2</sub>SiBe<sub>2</sub>O<sub>7</sub>*, Harry Diamond Laboratories, HDL-TR-1766 (August 1976).

Schmidt, P. C., A. Weiss, and T. P. Das, *Effect of crystal-fields and self-consistency on dipole and quadrupole polarizabilities of closed shell ions*, Phys. Rev. **B19** (1979), 5525.

Tessman, J. R., A. H. Kahn, and W. Shockley, Phys. Rev. **92** (1953), 890.

### A-3. $\text{Bi}_4\text{X}_3\text{O}_{12}$ ( $\text{X} = \text{Si}, \text{Ge}$ )

#### A-3.1 Crystallographic Data on $\text{Bi}_4\text{X}_3\text{O}_{12}$

Cubic  $T_d^6 (I\bar{4}3d)$ , 220,  $Z = 4$

Ion	Site	Symmetry	$x^a$	$y$	$z$	$q$	$\alpha (\text{\AA}^3)$
Bi	16c	$C_3$	$x$	$x$	$x$	3	$2.23^a$
X	12a	$S_4$	0	1/4	3/8	4	$\alpha_x^a$
O	48e	$C_1$	$x$	$y$	$z$	-4	$1.349^b$

<sup>a</sup>Fraga et al (1976),  $\alpha_{\text{Si}} = 0.03 (\text{\AA}^3)$ ,  $\alpha_{\text{Ge}} = 0.12 (\text{\AA}^3)$ .

<sup>b</sup>Schmidt et al (1979).

#### A-3.2 X-Ray Data on $\text{Bi}_4\text{X}_3\text{O}_{12}$

X	$a$	$x_{\text{Bi}}$	$x_0$	$y_0$	$z_0$
Si	10.300 <sup>a</sup>	0.0857	0.0607	0.1335	0.2875
Ge	10.513 <sup>b</sup>	0.0876	0.0689	0.1277	0.2875

<sup>a</sup>Wyckoff (1968), Vol. 4.

<sup>b</sup>Fisher and Waldner (1982).

#### A-3.3 Crystal-Field Components, $A_{kq} (\text{cm}^{-1}/\text{\AA}^k)$ , for the Si Site ( $S_4$ ) of $\text{Bi}_4\text{Si}_3\text{O}_{12}$

$A_{kq}$	Monopole	Self-induced	Dipole	Total
$A_{20}$	-4,423	1,103	11,851	8,531
$\text{Re}A_{32}$	39,156	-15,088	35,345	59,414
$\text{Im}A_{32}$	55,809	-21,662	43,774	77,921
$A_{40}$	-29,886	19,317	-36,749	-47,317
$\text{Re}A_{44}$	-7,631	4,551	-3,239	-6,319
$\text{Im}A_{44}$	18,425	-12,613	15,697	21,509
$\text{Re}A_{52}$	-2,043	1,506	2,727	2,190
$\text{Im}A_{52}$	-2,628	2,150	3,917	3,439
$ A_{44} $	19,943	—	—	22,418

### A-3.4 Crystal-Field Components, $A_{kq}$ ( $\text{cm}^{-1}/\text{\AA}^k$ ), for the Ge Site ( $S_4$ ) of $\text{Bi}_4\text{Ge}_3\text{O}_{12}$

$A_{kq}$	Monopole	Self-induced	Dipole	Total
$A_{20}$	-9,149	1,563	7,098	-488.4
$\text{Re}A_{32}$	26,496	-8,445	21,045	39,096
$\text{Im}A_{32}$	43,745	-14,017	29,123	58,851
$A_{40}$	-19,352	10,242	-22,771	-31,881
$\text{Re}A_{44}$	-7,790	3,872	-2,562	-6,480
$\text{Im}A_{44}$	13,318	-7,527	8,938	14,729
$\text{Re}A_{52}$	-2,366	1,523	1,117	274.0
$\text{Im}A_{52}$	-3,549	2,507	1,981	938.8
$ A_{44} $	15,429	—	—	16,091

### A-3.5 $\text{Bi}_4\text{X}_3\text{O}_{12}$ References

Allegretti, F., D. Borgia, R. Riva, F. DeNotaristefani, and S. Pizzini, *Growth of BGO single crystals using a directional solidification technique*, J. Cryst. Growth **94** (1989), 373.

Bochkova, T. M., E. G. Valyashko, V. A. Smirnov, and S. A. Flerova, *Optical spectra of  $\text{Bi}_4\text{Ge}_3\text{O}_{12}$  - Nd and  $\text{Bi}_4\text{Si}_3\text{O}_{12}$  - Nd crystals*, J. Appl. Spectrosc. (USSR) **30** (1979), 102.

Bortfeld, D. P., and H. Meier, *Refractive indices and electro-optic coefficients of the eulites  $\text{Bi}_4\text{Ge}_3\text{O}_{12}$  and  $\text{Bi}_4\text{Si}_3\text{O}_{12}$* , J. Appl. Phys. **43** (1972), 5110.

Chesler, R. B. and J. E. Geusic, *Laser handbook*, Vol.1, North Holland (1972), 362.

Cho, Z. H., O. Nalcioglu, and M. R. Farukhi, *Analysis of a cylindrical hybrid position camera with bismuth germanate (BGO) scintillation crystals*, IEEE NS25 (1978), 952.

Couzi, M., J. R. Vignalou, and G. Boulon, *Infrared and Raman study of the optical phonons in  $\text{Bi}_4\text{Ge}_3\text{O}_{12}$  single crystals*, Solid State Commun. **20** (1976), 461.

Fischer, P. and F. Waldner, *Comparison of neutron diffraction and EPR results on the cubic crystal structures of piezoelectric  $\text{Bi}_4\text{Y}_3\text{O}_{12}$  ( $Y = \text{Ge}, \text{Si}$ )*, Solid State Commun. **44** (1982), 657.

Gusev, V. A., and S. A. Petrov, *Photoluminescence of  $\text{Bi}_4\text{Ge}_3\text{O}_{12}$  single crystals*, Optoelectron. Instrum. Data Process. (5) (1988), 15.

Gusev, V. A., S. I. Demenko, and S. A. Petrov, *Photoconductivity of  $\text{Bi}_4\text{Ge}_3\text{O}_{12}$  single crystals*, Optoelectron. Instrum. Data Process. (32) (1988), 32.

Ivanov, V. Y., A. V. Kruzhalov, V. A. Pustovarou, and V. L. Petrov, *Electron excitation and luminescence in  $\text{Bi}_4\text{Ge}_3\text{O}_{12}$  and  $\text{Bi}_4\text{Si}_3\text{O}_{12}$  crystals*, Nucl. Instrum. Methods Phys. Res. A261 (1987), 150.

Johnson, L. F. and A. A. Ballman, *Coherent emission from rare earth ions in electro-optic crystals*, J. Appl. Phys. 40 (1969), 297.

Kaminiskii, A. A., S. E. Sarkisov, T. I. Butaeva, G. A. Denisenko, B. Hermoneit, J. Bohm, W. Grasskreutz, and D. Schultz, *Growth, spectroscopy, and stimulated emission of cubic  $\text{Bi}_4\text{Ge}_3\text{O}_{12}$  crystals doped with  $\text{Dy}^{3+}$ ,  $\text{Ho}^{3+}$ ,  $\text{Er}^{3+}$ ,  $\text{Tm}^{3+}$ , or  $\text{Yb}^{3+}$  ions*, Phys. Status Solidi (a)56 (1979), 725.

Kaminskii, A. A., D. Schultz, B. Hermoneit, S. Sarkisov, L. Li, J. Bohm, P. Reiche, R. Ehlert, A. A. Mayer, V. A. Lomonov, and V. A. Balashov, *Spectroscopic properties and stimulated emission in the  $^4F_{3/2} \rightarrow ^4I_{11/2}$  and  $^4F_{3/2} \rightarrow ^4I_{13/2}$  transitions of  $\text{Nd}^{3+}$  ions from cubic  $\text{Bi}_4\text{Ge}_3\text{O}_{12}$  crystals*, Phys. Status Solidi (a)33 (1976), 737.

Liebertz, J., *Einkristallzüchtung von wismutgermanat ( $\text{Bi}_4(\text{GeO}_4)_3$ )*, J. Cryst. Growth 5 (1969), 150.

Neikirk, D. P. and R. C. Powell, *Laser time-resolved spectroscopy studies of host-sensitized energy transfer in  $\text{Bi}_4\text{Ge}_3\text{O}_{12}$ :  $\text{Er}^{3+}$  crystals*, J. Lumin. 20 (1976), 261.

Nitsche, R., *Crystal growth and electro-optic effect of bismuth germanate,  $\text{Bi}_4(\text{GeO}_4)_3$* , J. Appl. Phys. 36 (1965), 2358.

Sarkisov, S. E., K. K. Pukhov, A. A. Kaminskii, A. G. Petrosyan, and T. I. Betaeva, *Manifestation of electron-phonon interaction in insulating crystals doped with  $\text{Pr}^{3+}$  ions*, Phys. Status Solidi (a)113 (1989), 193.

Segal, D. J., R. P. Santoro, and R. E. Newnham, *Neutron-diffraction study of  $\text{Bi}_4\text{Si}_3\text{O}_{12}$* , Z. Kristallogr. 123 (1966), 73.

Schmidt, P. C., A. Weiss, and T. P. Das, *Effect of crystal-fields and self-consistency on dipole and quadrupole polarizabilities of closed shell ions*, Phys. Rev. B19 (1979), 5525.

Schweppe, H., *Electromechanical properties of bismuth germanate  $\text{Bi}_4(\text{GeO}_4)_3$* , IEEE Trans. Sonics Ultrason. (October 1969), 219.

Von Philipsborn, H., *Croissance d'eulytine  $\text{Bi}_4\text{Si}_3\text{O}_{12}$  et des composés substitués  $\text{Bi}_4\text{Ge}_3\text{O}_{12}$  par la méthode czochrolski*, J. Cryst. Growth 11 (1971), 348.

Weber, M. J. and R. R. Monchamp, *Luminescence of  $\text{Bi}_4\text{Ge}_3\text{O}_{12}$ : Spectral and decay properties*, J. Appl. Phys. 44 (1973), 5495.

## Appendix A

### A-4. $\text{Bi}_{12}\text{XO}_{20}$ ( $\text{X} = \text{Ge}, \text{Si}$ )

#### A-4.1 Crystallographic Data on $\text{Bi}_{12}\text{XO}_{20}$

Cubic  $T^3$  ( $I23$ ), 197,  $Z = 2$

Ion	Site	Symmetry	$x$	$y$	$z$	$q$	$\alpha$ ( $\text{\AA}^3$ )
Bi	$24f$	$C_1$	$x$	$y$	$z$	3	$2.23^a$
X	$2a$	$T$	0	0	0	4	$\alpha_x^a$
O <sub>1</sub>	$24f$	$C_1$	$x$	$y$	$z$	-2	$1.349^b$
O <sub>2</sub>	$8c$	$C_3$	$x$	$x$	$x$	-2	1.349
O <sub>3</sub>	$8c$	$C_3$	$x$	$x$	$x$	-2	1.349

<sup>a</sup>Fraga et al (1976).

<sup>b</sup>Schmidt et al (1979).

#### A-4.2 X-Ray Data on $\text{Bi}_{12}\text{XO}_{20}$

X	$a$	$x_{\text{Bi}}$	$y_{\text{Bi}}$	$z_{\text{Bi}}$	$x_{\text{O1}}$	$y_{\text{O1}}$	$z_{\text{O1}}$
Ge	$10.1455^a$	0.82409	0.68158	0.98433	0.8655	0.7477	0.5145
Si	$10.10433^b$	0.17564	0.31741	0.01592	0.1348	0.2523	0.4858

X	$x_{\text{O2}}$	$x_{\text{O3}}$	$\alpha_x$ ( $\text{\AA}^3$ ) <sup>c</sup>
Ge	0.8019	0.0977	0.12
Si	0.1950	0.9059	0.03

<sup>a</sup>Abrahams et al (1967).

<sup>b</sup>Abrahams et al (1979).

<sup>c</sup>Fraga et al (1976).

#### A-4.3 Crystal-Field Components, $A_{kq}$ ( $\text{cm}^{-1}/\text{\AA}^k$ ), for the Ge Site of $\text{Bi}_{12}\text{GeO}_{20}$ <sup>a</sup>

$A_{kq}$	Monopole	Self-induced	Dipole	Total
$A_{32}$	$i54,550$	$-i17,875$	$i50,303$	86,978
$A_{40}$	$-25,350$	13,407	$-26,296$	$-38,240$
$A_{60}$	4,766.4	$-5,328.1$	7,328.2	6,766.5
$A_{62}$	163.03	17.182	$-35.978$	144.24
$A_{72}$	$-i3,901.1$	$i5,796.2$	$-i6,963.0$	$-i5,067.9$

$$^a A_{44} = \sqrt{\frac{5}{14}} A_{40}, A_{64} = -\sqrt{\frac{7}{2}} A_{60}, A_{66} = -\sqrt{\frac{5}{11}} A_{62}, \text{ and } A_{76} = \sqrt{\frac{11}{13}} A_{72}.$$



#### A-4.4 Crystal-Field Components, $A_{kq}$ ( $\text{cm}^{-1}/\text{\AA}^k$ ), for the Si Site ( $T$ ) of $\text{Bi}_{12}\text{SiO}_{20}$ <sup>a</sup>

$A_{kq}$	Monopole	Self-induced	Dipole	Total
$A_{32}$	-i64,514	i23,949	-i64,411	-i104,976
$A_{40}$	-31,029	18,698	-35,224	-47,555
$A_{60}$	6,361.3	-8,079.2	10,613	8,895.5
$A_{62}$	170.64	18.406	-35.384	153.67
$A_{72}$	i5,446.7	-i9,160.9	i10,524	i6,810.0

$$^a A_{44} = \sqrt{\frac{5}{14}} A_{40}, A_{64} = -\sqrt{\frac{7}{2}} A_{60}, A_{66} = -\sqrt{\frac{5}{11}} A_{62}, \text{ and } A_{76} = \sqrt{\frac{11}{13}} A_{72}.$$

#### A-4.5 $\text{Bi}_{12}\text{XO}_{20}$ ( $\text{X} = \text{Ge}, \text{Si}$ ) References

Abrahams, S. C., J. L. Bernstein, and C. Svensson, *Crystal structure and absolute piezoelectric  $d_{14}$  coefficient in laevorotatory  $\text{Bi}_{12}\text{SiO}_{20}$* , J. Chem. Phys. **71** (1979), 788.

Abrahams, S. C., P. B. Jamieson, and J. L. Bernstein, *Crystal structure of piezoelectric bismuth germanium oxide  $\text{Bi}_{12}\text{GeO}_{20}$* , J. Chem. Phys. **47** (1967), 4034.

Barbonas, G., D. Senulienė, A. Sileika, E. I. Leonov, and V. M. Orlov, *Induced circular dichroism in bismuth silicate ( $\text{Bi}_{12}\text{SiO}_{12}$ ) crystals doped with Cr*, Leit. Fiz. Rinkiny **27** (1987), 318 (in Russian; English abstract).

Denagbe, S., M. Martin, M. Schvocrer, F. Marsaud, J. C. Launay, and P. Hagenmuller, *Thermally stimulated luminescence in undoped and doped  $\text{Bi}_{12}\text{GeO}_{20}$  single crystals*, J. Phys. Chem. Solids **51** (1990), 171.

Fraga, S., K. M. Saxena, and J. Karwowski, *Physical science data: 5. Handbook of atomic data*, Elsevier, New York (1976).

Nikolova, E. G. and M. M. Gospodinov, *The Twyman effect in crystals of  $\text{Bi}_4\text{Ge}_3\text{O}_{12}$  and  $\text{Bi}_{12}\text{SiO}_{20}$* , J. Mater. Sci. Lett. **8** (1989), 309.

Schmidt, P. C., A. Weiss, and T. P. Das, *Effect of crystal-fields and self-consistency on dipole and quadrupole polarizabilities of closed shell ions*, Phys. Rev. **B19** (1979), 5525.

Senulienė, D., G. Babonas, A. Sileika, E. Leonovas, and V. Orlovas, *An investigation of absorption and circular dichroism spectra in  $\text{Bi}_{12}\text{SiO}_{20}$  crystals doped with Ni, Co, and Mn*, Sov. Phys. Coll. **27** No. 1 (1987), 55 (English translation).

Wardzynski, W., M. Baran, and H. Szymczak, *Electron paramagnetic resonance of  $\text{Fe}^{3+}$  in bismuth germanium oxide single crystals*, Physica **111B** (1981), 47.

Wardzynski, W., T. Lukasiewicz, and J. Zmija, *Reversible photochromic effects in doped single crystals of bismuth germinate ( $\text{Bi}_{12}\text{GeO}_{20}$ ) and bismuth silicon oxide ( $\text{Bi}_{12}\text{SiO}_{20}$ )*, Opt. Commun. **30** (1979), 203.

Wardzynski, W., and H. Szymczak, *The center of orthorhombic symmetry in chromium doped  $\text{Bi}_{12}\text{GeO}_{20}$  and  $\text{Bi}_{12}\text{SiO}_{20}$  single crystals*, J. Phys. Chem. Solids **45** (1984), 887.

Wardzynski, W., H. Szymczak, K. Pataj, T. Lukasiewicz, and J. Zmija, *Light induced charge transfer in Cr doped  $\text{Bi}_{12}\text{GeO}_{20}$  and  $\text{Bi}_{12}\text{SiO}_{20}$  single crystals*, J. Phys. Chem. Solids **43** (1982), 767.

Weber, M. J., ed., *CRC handbook of laser science and technology, Vol. IV. Optical materials, Part 2: Properties*, CRC Press, Boca Raton, FL (1986). Also, Vol. V. *Optical materials, Part 3: Applications, coatings, and fabrication*, CRC Press (1987).

Wojdowski, W., T. Lukasiewicz, W. Nazarewicz, and J. Zmija, *Infrared studies of lattice vibrations in  $\text{Bi}_{12}\text{GeO}_{20}$  and  $\text{Bi}_{12}\text{SiO}_{20}$  crystals*, Phys. Status Solidi (b) **94** (1979), 649.

## DISTRIBUTION

Administrator  
Defense Technical Information Center  
Attn DTIC-DDA (2 copies)  
Cameron Station, Bulding 5  
Alexandria, VA 22304-6145

Director  
Night Vision & Electro-Optics Laboratory  
Attn Technical Library  
Attn R. Buser  
Attn A. Pinto  
Attn J. Hebersat  
Attn L. Merkel  
Attn R. Rhode  
Attn W. Tressel  
Ft Belvoir, VA 22060

Director  
Defense Advanced Research Projects Agency  
Attn J. Friebele  
1400 Wilson Blvd  
Arlington, VA 22290

Director  
Defense Nuclear Agency  
Attn Tech Library  
Washington, DC 20305

Under Secretary of Defense Res &  
Engineering  
Attn Tech Library, 3C128  
Washington, DC 20301

Office of the Deputy Chief of Staff, for  
R&D Center  
Department of the Army  
Attn DAMA-ARZ-B,  
I. R. Hershner  
Washington, DC 20310

Commander  
US Army Armament Munitions &  
Chemical Command (AMCCOM)  
R&D Center  
Attn DRDAR-TSS, STINFO Div  
Dover, NJ 07801

Commander  
Atmospheric Sciences Laboratory  
Attn Technical Library  
White Sands Missile Range, NM 88002

Director  
US Army Ballistic Research Laboratory  
Attn SLCBR-DD-T (STINFO)  
Aberdeen Proving Ground, MD 21005

Director  
US Army Electronics Warfare Laboratory  
Attn J. Charlton  
Attn AMSEL-DD  
Ft Monmouth, NJ 07703

Commanding Officer  
USA Foreign Science & Technology  
Center  
Attn AIAST-BS, Basic Science Div  
Federal Office Building  
Charlottesville, VA 22901

Commander  
US Army Materials & Mechanics  
Research Center  
Attn SLCMT-TL, Tech Library  
Watertown, MA 02172

US Army Materiel Systems Analysis  
Activity  
Attn AMXSU-MP, Library  
Aberdeen Proving Ground, MD 21005

Commander  
US Army Missile & Munitions  
Center & School  
Attn ATSK-CTD-F  
Attn AMSMI-TB, Redstone Sci Info  
Center  
Redstone Arsenal, AL 35809

Commander  
US Army Research Office Durham  
Attn M. Stosio  
Attn M. Ciftan  
Attn R. Guenther  
Research Triangle Park, NC 27709

Commander  
USA Resch & Std Gp (Europe)  
Attn Chief, Physics & Math Branch  
FPO, New York 09510

## DISTRIBUTION

Commander  
US Army Test & Evaluation Command  
Attn D. H. Sliney  
Attn Tech Library  
Aberdeen Proving Ground, MD 21005

Commander  
US Army Troop Support Command  
Attn STRNC-RTL, Tech Library  
Natick, MA 01762

Office of Naval Research  
Attn J. Murday  
Arlington, VA 22217

Director  
Naval Research Laboratory  
Attn Code 2620, Tech Library Br  
Attn G. Quarles  
Attn G. Kintz  
Attn A. Rosenbaum  
Attn G. Risenblatt  
Attn Code 5554, F. Bartoli  
Attn Code 6540, S. R. Bowman  
Attn Code 5554, L. Esterowitz  
Attn Code 5554, R. E. Allen  
Washington, DC 20375

Commander  
Naval Weapons Center  
Attn Code 3854, R. Schwartz  
Attn Code 3854, M. Hills  
Attn Code 3854, M. Nadler  
Attn Code 385, R. L. Atkins  
Attn DOCE 343, Technical Information  
Department  
China Lake, CA 93555

Air Force Office of Scientific Research  
Attn Major H. V. Winsor, USAF  
Bolling AFB  
Washington, DC 20332

Department of Commerce  
National Bureau of Standards  
Attn Library  
Washington, DC 20234

Nasa Langley Research Center  
Attn N. P. Barnes  
Attn G. Armagan

Nasa Langley Research Center (cont'd)  
Attn P. Cross  
Attn D. Getteny  
Attn J. Barnes  
Attn E. Filer  
Attn C. Bair  
Attn N. Bounchrishani  
Hampton, VA 23665

Director  
Advisory Group on Electron Devices  
Attn Sectry, Working Group D  
201 Varick Street  
New York, NY 10013

Aerospace Corporation  
Attn M. Birnbaum  
Attn N. C. Chang  
Attn T. S. Rose  
PO Box 92957  
Los Angeles, CA 90009

Allied Signal, Inc.  
Advanced Application Dept  
Attn A. Budgor  
31717 La Tienda Drive  
Westlake Village, CA 91362

Allied Signal Inc  
Attn Y. Band  
Attn R. Morris  
POB 1021R  
Morristown, NJ 07960

Ames Laboratory Dow  
Iowa State University  
Attn K. A. Gschneidner, Jr (2 copies)  
Ames, IA 50011

Argonne National Laboratory  
Attn W. T. Carnall  
9700 South Cass Avenue  
Argonne, IL 60439

Booz Allen and Hamilton  
Attn W. Drozdowski  
4330 East West Highway  
Bethesda, MD 20814

## DISTRIBUTION

Brimrose Corp of America  
Attn R. G. Rosemeier  
7527 Belair Road  
Baltimore, MD 21236

Draper Lab  
Attn F. Hakimi, MS 53555 Tech Sq  
Cambridge, MA 02139

Engineering Societies Library  
Attn Acquisitions Dept  
345 East 47th Street  
New York, NY 10017

Fibertech Inc  
Attn H. R. Verdin (3 copies)  
510-A Herdon Pkwy  
Herdon, VA 22070

General Dynamics  
Attn R. J. Blair  
5452 Oberlin Drive  
San Diego, CA 92121

Hughes Aircraft Company  
Attn D. Sumida  
3011 Malibu Canyon Rd  
Malibu, CA 90265

IBM Research Division  
Almaden Research Center  
Attn R. M. Macfarlane,  
Mail Stop K32 802(d)  
650 Harry Road  
San Jose, CA 95120

Director  
Lawrence Radiation Laboratory  
Attn M. J. Weber  
Attn H. A. Koehler  
Attn W. Krupke  
Livermore, CA 94550

LTV  
Attn M. Kock (WT-50)  
PO Box 650003  
Dallas, TX 75265

Martin Marietta  
Attn F. Crowne  
Attn J. Little

Martin Marietta (cont'd)  
Attn T. Worchesky  
1450 South Rolling Road  
Baltimore, MD 21227

McDonnell Douglass Electronic Systems  
Company  
Dept Y440, Bldg. 101, Lev. 2 Rm/Pt B54  
Attn D. M. Andrauskas, MS-2066267  
PO Box 516  
St Louis, MO 63166

MIT Lincoln Lab  
Attn B. Aull  
PO Box 73  
Lexington, MA 02173

Department of Mechanical, Industrial, &  
Aerospace Engineering  
Attn S. Temkin  
PO Box 909  
Piscataway, NJ 08854

National Oceanic & Atmospheric Adm  
Environmental Research Labs  
Attn Library, R-51, Tech Rpts  
Boulder, CA 80302

Oak Ridge National Laboratory  
Attn R. G. Haire  
Oak Ridge, TN 37839

W. J. Schafer Assoc  
Attn J. W. Collins  
321 Billerica Road  
Chelmsford, MA 01824

Science Applications, International Corp  
Attn T. Allik  
1710 Goodridge Drive  
McLean, VA 22102

Shwartz Electro-Optic, Inc.  
Attn G. A. Rines  
45 Winthrop Street  
Concord, MA 01742

Teledyne Brown Engineering  
Cummings Research Park  
Attn M. L. Price, MS-44  
Huntsville, AL 35807

## DISTRIBUTION

Union Carbide Corp  
Attn M. R. Kokta  
50 South 32nd Street  
Washougal, WA 98671

Arizona State University  
Dept of Chemistry  
Attn L Eyring  
Tempe, AZ 85281

University of Southern California  
Attn Dr. M. Birnbaum  
Denney Research Bldg, University Park  
Los Angeles, CA 90089

Carnegie Mellon University  
Schenley Park  
Attn Physics & EE, J. O. Artman  
Pittsburgh, PA 15213

Colorado State University  
Physics Department  
Attn S. Kern  
Ft Collins, CO 80523

University of Connecticut  
Department of Physics  
Attn R. H. Bartram  
Storrs, CT 06269

University of South Florida  
Physics Dept  
Attn R. Chang  
Attn Sengupta  
Tampa, FL 33620

Howard University  
Physics Department  
Attn Prof. V. Kushamaha  
25 Bryant St., N.W.  
Washington, DC 20059

Johns Hopkins University  
Dept of Physics  
Attn B. R. Judd  
Baltimore, MD 21218

Kalamazoo College  
Dept of Physics  
Attn K. Rajnak  
Kalamazoo, MI 49007

Massachusetts Institute of Technology  
Crystal Physics Laboratory  
Attn H. P. Jenssen  
Cambridge, MA 02139

Massachusetts Institute of Technology  
Attn V. Bagnato  
77 Mass. Ave, Room 26-251  
Cambridge, MA 02139

University of Minnesota, Duluth  
Department of Chemistry  
Attn L. C. Thompson  
Duluth, MN 55812

Oklahoma State University  
Dept of Physics  
Attn R. C. Powell  
Stillwater, OK 74078

Pennsylvania State University  
Materials Research Laboratory  
Attn W. B. White  
University Park, PA 16802

Princeton University  
Department of Chemistry  
Attn D. S. McClure  
Princeton, NJ 08544

San Jose State University  
Department of Physics  
Attn J. B. Gruber  
San Jose, CA 95192

Seaton Hall University  
Chemistry Department  
Attn H. Brittain  
South Orange, NJ 07099

University of Virginia  
Dept of Chemistry  
Attn F. S. Richardson (2 copies)  
Charlottesville, VA 22901

University of Wisconsin  
Chemistry Department  
Attn J. Wright  
Attn B. Tissue  
Madison, WI 53706

## DISTRIBUTION

US Army Laboratory Command  
Attn Technical Director, AMSLC-TD

Installation Support Activity  
Attn Legal Office, SICIS-CC

USAISC  
Attn Administrative Services, AMSLC-IM-TS  
Attn Technical Publishing Branch,  
AMSLC-IM-VP (2 copies)

Harry Diamond Laboratories  
Attn Division Directors  
Attn Library, SLCHD-TL (3 copies)  
Attn Library, SLCHD-TL (Woodbridge)  
Attn Chief, SLCHD-NW-CS  
Attn Chief, SLCHD-NW-E  
Attn Chief, SLCHD-NW-EH  
Attn Chief, SLCHD-NW-EP  
Attn Chief, SLCHD-NW-ES  
Attn Chief, SLCHD-NW-P  
Attn Chief, SLCHD-NW-R  
Attn Chief, SLCHD-NW-RP  
Attn Chief, SLCHD-NW-RS  
Attn Chief, SLCHD-NW-TN  
Attn Chief, SLCHD-NW-TS  
Attn Chief, SLCHD-PO  
Attn Chief, SLCHD-ST-C  
Attn Chief, SLCHD-ST-RP  
Attn Chief, SLCHD-ST-RS  
Attn Chief, SLCHD-TT

Harry Diamond Laboratories (cont'd)  
Attn B. Willis, SLCHD-TA-ET  
Attn B. Zabloudowski, SLCHD-TA-ET  
Attn C. S. Kenyon, SLHD-NW-EP  
Attn J. R. Miletta, SLCHD-NW-EP  
Attn F. B. McLean, SLCHD-ST-MW  
Attn L. Libelo, SLCHD-ST-MW  
Attn A. A. Bencivenga, SLCHD-ST-SP  
Attn J. Sattler, SLCHD-CS  
Attn J. Nemarich, SLCHD-ST-CB  
Attn B. Weber, SLCHD-ST-CB  
Attn T. Bahder, SLCHD-ST-AP  
Attn J. Bradshaw, SLCHD-ST-AP  
Attn J. Bruno, SLCHD-ST-AP  
Attn E. Harris, SLCHD-ST-AP  
Attn R. Leavitt, SLCHD-ST-AP  
Attn J. Pham, SLCHD-ST-AP  
Attn G. Simonis, SLCHD-ST-AP  
Attn M. Stead, SLCHD-ST-AP  
Attn J. Stellato, SLCHD-ST-AP  
Attn S. Stevens, SLCHD-ST-AP  
Attn R. Tober, SLCHD-ST-AP  
Attn M. Tobin, SLCHD-ST-AP  
Attn G. Turner, SLCHD-ST-AP  
Attn D. Wortman, SLCHD-ST-AP  
Attn C. Garvin, SLCHD-ST-SS  
Attn J. Goff, SLCHD-RT-RB  
Attn C. Morrison, SLCHD-ST-AP  
(10 copies)